

EXTENSIONS OF LIE BRACKETS.

OLIVIER BRAHIC

ABSTRACT. We provide a framework for extensions of Lie algebroids, including non-abelian extensions and Lie algebroids over different bases. Our approach involves Ehresmann connections, which allows straight generalizations of classical constructions. We exhibit a filtration in cohomology and explain the associated spectral sequence. We also give a description of the groupoid integrating an extension in case a complete connection exists. The integrability is also studied.

INTRODUCTION

The concept of Lie algebroid proved to be relevant in diverse fields of geometry, with interesting applications in physics as well. However, despite intensive studies, the notion of extension was never systematically investigated. It is the aim of this paper to fill this gap. We study extensions of Lie algebroids from a general perspective, the interest is to get a unified treatment for a large variety of situations.

The basic setting will be a surjective Lie algebroid morphism $\pi : A_E \rightarrow A_B$ covering a surjective submersion $p : E \rightarrow B$ where A_E and A_B are Lie algebroids over E and B , respectively. In this situation, we denote $\mathcal{K}$ the kernel of π . It is a Lie algebroid over E and we obtain this way A_E as an extension of A_B by $\mathcal{K}$:

$$\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B.$$

In Section 1, we recall basic notion concerning Lie algebroids and their extensions. Then we introduce in Section 2 the notion of Ehresmann connection for an extension: it is given by a sub-vector bundle H complementary to $\mathcal{K}$ in A_E :

$$A_E = \mathcal{K} \oplus H.$$

Such a connection always exists and we can construct the usual horizontal lifting, it is a $C^\infty(B)$ -linear application $h : \Gamma(A_B) \rightarrow \Gamma(H)$. From these notions, we derive a couple $(\mathcal{D}, \omega)$ where:

- $\mathcal{D} : \Gamma(A_B) \rightarrow \text{Der}(\mathcal{K})$ is a $C^\infty(B)$ -linear map with values in derivations of $\mathcal{K}$, such that the symbol $s_{\mathcal{D}_\alpha}$ of $\mathcal{D}_\alpha$ p_* -projects onto $\sharp_B(\alpha)$ for any $\alpha \in \Gamma(A_B)$;
- ω is a 2-form on A_B with values in sections of $\mathcal{K}$: $\omega \in \Omega^2(A_B) \otimes \Gamma(\mathcal{K})$.

Partially supported by the Fundação para a Ciência e a Tecnologia through the Program POCI 2010/FEDER and by the Projects POCI/MAT/57888/2004 and POCI/MAT/55958/2004.

In order to define an extension, the couple $(\mathcal{D}, \omega)$ is submitted to the following compatibility conditions:

$$\begin{aligned}\partial_H \omega &= 0, \\ \text{Curv}_{\mathcal{D}} &= \text{ad}^{\mathcal{K}} \circ \omega.\end{aligned}$$

Here, $\partial_H : \Omega^k(A_B) \otimes \Gamma(\Lambda^l \mathcal{K}) \rightarrow \Omega^{k+1}(A_B) \otimes \Gamma(\Lambda^l \mathcal{K})$ is a covariant derivative associated to $\mathcal{D}$ by usual formulas and $\text{Curv}_{\mathcal{D}}(\alpha, \beta) := [\mathcal{D}\alpha, \mathcal{D}\beta] - \mathcal{D}_{[\alpha, \beta]}$.

The description of extensions we get this way lies in-between usual treatments for both Lie algebras and fibrations:

Theorem. *Let $p : E \rightarrow B$ be a submersion, $\mathcal{K} \rightarrow E$ a Lie algebroid whose characteristic foliation lies in Vert and $A_B \rightarrow B$ a Lie algebroid.*

Extensions of A_B by $\mathcal{K}$ covering p are (up to equivalence) classified by equivalence classes of admissible couples $(\mathcal{D}, \omega)$ under the relation $(\mathcal{D}, \omega) \sim (\mathcal{D}', \omega')$ if there exists $\Delta \in \Omega^1(A_B) \otimes \Gamma(\mathcal{K})$ such that

$$\begin{aligned}\mathcal{D}' &= \mathcal{D} + \text{ad}^{\mathcal{K}} \circ \Delta, \\ \omega' &= \omega + \partial_H \Delta + [\Delta \wedge \Delta]_{\mathcal{K}},\end{aligned}$$

where $[\Delta \wedge \Delta]_{\mathcal{K}} \in \Omega^2(A_B) \otimes \Gamma(\mathcal{K})$ is defined by:

$$[\Delta \wedge \Delta]_{\mathcal{K}}(\alpha, \beta) = [\Delta(\alpha), \Delta(\beta)]_{\mathcal{K}}.$$

Besides, we explain that given a connection, parallel transport along an A_B -path $a \in P(A_B)$ is well defined (see subsection 2.1) and, in the case of a complete connection, we obtain the usual notion of holonomy along a as a Lie algebroid morphism

$$\mathcal{K}|_{E_{\gamma(0)}} \xrightarrow{\Phi_a} \mathcal{K}|_{E_{\gamma(1)}}.$$

Here, $\gamma : I \rightarrow B$ is the base path of a . In general, Φ does not factor through A_B -homotopies unless $\text{Curv}_{\mathcal{D}}$ vanishes. Note however the relation $\text{Curv}_{\mathcal{D}} = \text{ad}^{\mathcal{K}} \circ \omega$ suggests that this can be measured by inner automorphisms of $\mathcal{K}$, this will play an important role. In any case, one can still think of Φ as a groupoid morphism $\Phi : P(A_B) \rightarrow \text{Gauge}(\mathcal{K})$ where

$$\text{Gauge}(\mathcal{K}) := \{\text{Lie algebroid morphisms } \Psi_{y,x} : \mathcal{K}|_{E_x} \rightarrow \mathcal{K}|_{E_y}\}$$

and $P(A_B)$ are both groupoids over B .

In Section 3 we look at the cohomology of an extension. There is a natural filtration, and the choice of a connection allows a rather nice description of the associated spectral sequence.

The Section 4 is devoted to the description of the Weinstein groupoid $\mathcal{G}(A_E)$. We obtain $\mathcal{G}(A_E)$ as a quotient of $P(A_B) \ltimes \mathcal{G}(\mathcal{K})$. Here $P(A_B)$ acts on $\mathcal{G}(\mathcal{K})$ by means of holonomy, and $P(A_B) \ltimes \mathcal{G}(\mathcal{K})$ corresponds to an associated groupoid over E . The precise statement is given by Theorem 4.5:

Theorem. *Consider a Lie algebroid extension $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ endowed with a Ehresmann connection $A_E = \mathcal{K} \oplus H$, which we assume to be complete.*

Then the topological source simply connected groupoid $\mathcal{G}(A_E)$ integrating A_E is naturally identified with the quotient

$$P(A_B) \ltimes_B \mathcal{G}(\mathcal{K}) / \sim,$$

where the equivalence relation is given by: $(a_0, g_0) \sim (a_1, g_1)$ if and only if there exists a A_B -homotopy $h_B = adt + bde : TI^2 \mapsto A_B$ between a_0 and a_1 such that:

$$g_1 \cdot g_0^{-1} = \partial(h_B, \mathbf{t}(g_0)).$$

Here, $\partial(h_B, x_0)$ is the element in $\mathcal{G}(\mathcal{K})$ represented by the $\mathcal{K}$ -path:

$$\epsilon \rightarrow \int_0^1 (\Phi_{0,s}^{a_B^\epsilon})_*(\omega(a, b)_{s,\epsilon}) ds \in \mathcal{K}_{\gamma_K^\epsilon},$$

where $\gamma_K^\epsilon := \phi_{a_B^\epsilon}^{-1} \circ \phi_{a_B^0}(x_0)$.

In general, the sequence of topological groupoids

$$1 \rightarrow \mathcal{G}(\mathcal{K}) \xrightarrow{\tilde{\iota}} \mathcal{G}(A_E) \xrightarrow{\tilde{\pi}} \mathcal{G}(A_B) \rightarrow 1.$$

obtained by integration of a Lie algebroid extension might not be exact. In order to measure this lack of exactness, we denote $\mathcal{M}$ the monodromy groupoid of the extension. It is defined as the kernel of $\tilde{\iota}$. In the case there exists a complete Ehresmann connection, we can build a connecting homomorphism whose image is precisely $\mathcal{M}$, similarly to what is well known for fibrations:

Theorem. *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a Lie algebroid extension that admits a complete Ehresmann connection. Then there exists a homomorphism*

$$\partial_2 : \pi_2(A_B) \times_B E \rightarrow \mathcal{G}(\mathcal{K}),$$

that makes the following sequence exact:

$$\cdots \rightarrow \pi_2(A_B) \times E \xrightarrow{\partial_2} \mathcal{G}(\mathcal{K}) \xrightarrow{\tilde{\iota}} \mathcal{G}(A_E) \twoheadrightarrow \mathcal{G}(A_B).$$

Here, $\pi_2(A_B)$ denote the second homotopy groups of A_B , as defined in the Appendix C. In the case A_B is integrable, it coincides with the union of all second homotopy groups of the source fibers, seen as a bundle of groups over B . In this work we do not try to put any topology on $\pi_2(A_B)$ whose role is essentially algebraic.

The interest of the monodromy groupoid $\mathcal{M}$ lies in the fact that it controls the integrability of A_E in the case of *clean* extensions. By *clean*, we mean by definition that the restrictions of π to the isotropy Lie algebras of A_E surject onto those of A_B . This implies that, pointwise, one has exact sequences of Lie algebras

$$(\ker \sharp_{\mathcal{K}})_x \hookrightarrow (\ker \sharp_E)_x \twoheadrightarrow (\ker \sharp_B)_{p(x)}.$$

In this case, we show that A_E is integrable provided A_B and $\mathcal{K}$ are, and if $\mathcal{M}$ is discrete near identities (see 4.17 for the precise definition).

Theorem. *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a clean Lie algebroid extension. Assume that it admits a complete Ehresmann connection and that both $\mathcal{K}$ and A_B are integrable Lie algebroids.*

Then A_E is integrable if $\mathcal{M}$ is discrete near identities in $\mathcal{G}(\mathcal{K})$.

Acknowledgments. I would like to thank Rui Loja Fernandes for the many useful discussions and suggestions that helped to improve this work.

CONTENTS

Introduction	1
1. Generalities on Lie algebroids	4
1.1. Derivations.	5
1.2. Extensions of Lie algebroids	6
2. Ehresmann Connections	7
2.1. Parallel transport	8
2.2. Admissible couples	9
2.3. Isotropies	10
2.4. Semi-direct products	11
2.5. Abelian Extensions	11
2.6. Examples	11
3. Cohomology	13
3.1. The spectral sequence.	13
3.2. Interpretation	17
3.3. An example of computation	18
4. Integration	19
4.1. Splitting A_E -paths and A_E -homotopies	19
4.2. The Monodromy Groupoid	24
4.3. Integrability	27
Appendix A. The Weinstein groupoid and A -homotopies	28
Appendix B. Fibered Lie groupoids	30
Appendix C. Spheres in a Lie algebroid	31
References	33

1. GENERALITIES ON LIE ALGEBROIDS

We first briefly recall basic definitions concerning Lie algebroids [16] and refer to [3], [13] for more details.

Definition 1.1. A **Lie algebroid** $A \rightarrow M$ is a vector bundle A over a smooth manifold M , together with vector bundle homomorphism $\sharp : A \rightarrow TM$ called the anchor, and a bracket $[\cdot, \cdot]_A$ defined on the space $\Gamma(A)$ of sections of A such that for any smooth function $f \in C^\infty(M)$ and any section $\alpha, \beta \in \Gamma(A)$ of A , the following conditions hold

- i) $[\alpha, f\beta]_A = f[\alpha, \beta]_A + \mathcal{L}_{\sharp(\alpha)}(f)\beta$;
- ii) $[\alpha, \beta]_A = -[\beta, \alpha]_A$;
- iii) $\sharp([\alpha, \beta]_A) = [\sharp(\alpha), \sharp(\beta)]_{TM}$;
- iv) $\oint_{\alpha, \beta, \gamma} [[\alpha, \beta]_A, \gamma]_A = 0$.

In other words, the bracket $[\cdot, \cdot]_A$ induces a structure of real Lie algebra on sections of A such that the Leibniz condition (i) holds and $\sharp$ is a Lie algebra homomorphism from $\Gamma(A)$ to $\mathfrak{X}(M) := \Gamma(TM)$.

The cohomology associated to a Lie algebroid is defined on the complex of cochains $C^k := \Gamma(\Lambda^k A^*)$ which we might also denote $\Omega^k(A)$ (or $\Omega^k(M)$ for short when $A = TM$). The differential operator is given by the usual

formula

$$(1) \quad \langle d_A c, \alpha_0 \dots \alpha_k \rangle := \sum_{i=0 \dots n} (-1)^i \mathcal{L}_{\sharp \alpha_i} \langle c, \alpha_0 \dots \hat{\alpha}_i \dots \alpha_k \rangle \\ + \sum_{0 \leq i < j \leq k} (-1)^{i+j} \langle c, [\alpha_i, \alpha_j]_A, \alpha_0 \dots \hat{\alpha}_i \dots \hat{\alpha}_j \dots \alpha_k \rangle.$$

Definition 1.2. Given two Lie algebroids $A_M \rightarrow M$ and $A_N \rightarrow N$, a **Lie algebroid homomorphism** $\Phi : A_M \rightarrow A_N$ over $\phi : M \rightarrow N$ is a vector bundle homomorphism Φ covering ϕ such that the map induced on forms:

$$\Phi^* : \Omega^k(A_N) \rightarrow \Omega^k(A_M),$$

defined by $\langle \Phi^*(u), a \rangle = \langle u_{\phi(m)}, \Phi(a) \rangle$, commutes with the differential:

$$d_{A_M} \circ \Phi^* = \Phi^* \circ d_{A_N}.$$

When ϕ is a diffeomorphism, we shall denote $\Phi_* : \Gamma(A) \rightarrow \Gamma(A)$ the application induced on sections, $\Phi_*(\alpha) := \Phi \circ \alpha \circ \phi^{-1}$. See for instance [12] for a detailed discussion on Lie algebroid morphisms.

Recall that the isotropy at a point $x \in M$ is defined as the kernel of the anchor, $\mathfrak{g}_x^A := \ker \sharp_x$. It is naturally endowed with a structure of Lie algebra. In general, the dimension of the isotropy algebra may vary from one point to another, so all isotropies do not fit in a smooth bundle of Lie algebras in general.

1.1. Derivations.

Definition 1.3. A **derivation** of a Lie algebroid $A \rightarrow M$ is an application

$$D : \Gamma(A) \rightarrow \Gamma(A),$$

together with a vector field $s_D \in \mathfrak{X}(M)$ (called the **symbol** of the derivation) satisfying for any smooth function $f \in C^\infty(M)$ and any two sections of A $\alpha, \beta \in \Gamma(A)$

$$D(f\alpha) = fD(\alpha) + s_D(f)\alpha, \\ D([\alpha, \beta]_A) = [D(\alpha), \beta]_A + [\alpha, D(\beta)]_A, \\ \sharp(D(\alpha)) = [s_D, \sharp(\alpha)]_{TM}.$$

A derivation is the infinitesimal version of a Lie algebroid automorphism: any derivation determines a vector field on A whose flow is a Lie algebroid automorphism *et vice versa* (see for instance the appendix in [6]). We will denote $\text{Der}(A)$ the space of all derivations, it has a natural structure of Lie algebra, whose bracket is given by the commutator $[D_1, D_2] = D_1 \circ D_2 - D_2 \circ D_1$.

Definition 1.4. An **inner derivation** is a derivation of the form $D(\alpha) = [\delta, \alpha]_A$ for some smooth section $\delta \in \Gamma(A)$, in which case we will write $D = \text{ad}_\delta^A$.

There are obvious notions of time-dependent derivations and inner derivations, corresponding to which we get time-dependent vector fields on A . In particular, given a time-dependent section $\alpha^t \in \Gamma(A)$, we shall denote $\psi_{t,s}^\alpha : A \rightarrow A$ the flow on A associated to $\text{ad}_{\alpha^t}^A$.

1.2. Extensions of Lie algebroids. The notion of extension we will use in this work is summarized below.

Definition 1.5. An **extension** of a Lie algebroid A_B is a surjective Lie algebroid morphism $\pi : A_E \rightarrow A_B$ covering a surjective submersion $p : E \rightarrow B$.

With this definition, $\mathcal{K} := \ker \pi$ is a smooth sub-vector bundle of A_E that, as proven below, turns out to be a Lie sub-algebroid of A_E . Thus, we obtain an exact sequence of Lie algebroids:

$$\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B.$$

We shall refer to A_E as an extension of A_B by $\mathcal{K}$. As generic notations, we will write $\sharp_E : A_E \rightarrow TE$, $\sharp_B : A_B \rightarrow TB$, $\sharp_{\mathcal{K}} : \mathcal{K} \rightarrow TE$ and $[\cdot, \cdot]_{A_E}$, $[\cdot, \cdot]_{A_B}$, $[\cdot, \cdot]_{\mathcal{K}}$ the respective anchors and brackets. The isotropy algebras will be denoted $\mathfrak{g}_x^E := \ker(\sharp_E)_x$, $\mathfrak{g}_b^B := \ker(\sharp_B)_b$, and $\mathfrak{g}_x^{\mathcal{K}} := \ker(\sharp_{\mathcal{K}})_x = \mathfrak{g}_x^E \cap \mathcal{K}$, for any $x \in E, b \in B$.

Remark 1.6. Note that the injection $\mathcal{K} \subset A_E$ does not determine π (not even p) so one should not think of A_B as a quotient of A_E by $\mathcal{K}$.

Definition 1.7. Given an extension $\pi : A_E \twoheadrightarrow A_B$, we will refer to a π -**projectable** section μ of A_E any section μ such that, for any form $\nu \in \Omega^1(A_B)$, the contraction $\langle \pi \circ \mu, \nu \rangle$ is a basic function on $E \rightarrow B$. This means that $\pi \circ \mu$ induces a well defined section of A_B that we will denote $\pi(\mu)$.

Lemma 1.8. Let $\mu \in \Gamma(A_E)$ be a π -projectable section, then $\sharp_E \circ \mu$ is p_* -projectable, moreover:

$$p_*(\sharp_E(\mu)) = \sharp_B(\pi(\mu)).$$

Proof. Recall the condition for π to be a Lie algebroid morphism: $d_{A_E} \circ \pi^* = \pi^* \circ d_{A_B}$ where $\pi^* : \Omega^k(A_B^*) \rightarrow \Omega^k(A_E)$ is naturally induced by π . For $k = 0$, this condition writes:

$$\begin{aligned} \langle d_{A_E} \circ \pi^*(f), \mu \rangle &= \langle \pi^* \circ d_{A_B}(f), \mu \rangle & (f \in C^\infty(B), \mu \in \Gamma(A_E)) \\ \iff \langle d(\pi^*f), \sharp_E(\mu) \rangle &= \langle d_A(f), \pi \circ \mu \rangle \\ \iff \langle d(f \circ p), \sharp_E(\mu) \rangle &= \langle df, \sharp_B \circ \pi \circ \mu \rangle \\ \iff \langle df \circ dp, \sharp_E(\mu) \rangle &= \langle df, \sharp_B \circ \pi \circ \mu \rangle, \end{aligned}$$

which is *à priori* an equality in $C^\infty(E)$. However, if we assume μ to be π -projectable, the right-hand term is in fact a basic function, which implies that $\sharp_E(\mu)$ p_* -projects onto $\sharp_B(\pi(\mu))$. $\square$

Lemma 1.9. Let $\mu, \nu \in \Gamma(A_E)$ be π -projectable sections, then $[\mu, \nu]_{A_E}$ is also projectable, moreover:

$$\pi([\mu, \nu]_{A_E}) = [\pi(\mu), \pi(\nu)]_{A_B}.$$

Proof. This is an immediate consequence of the condition $d_{A_E} \circ \pi^* = \pi^* \circ d_{A_B}$ $\pi^* : \Omega^k(A_B) \rightarrow \Omega^k(A_E)$ for $k = 1$ and of the preceding lemma. $\square$

As a consequence of the two last lemmas, we can state the following.

Proposition 1.10. *Let $\pi : A_E \rightarrow A_B$ be a Lie algebroid extension. Then $\mathcal{K} := \ker \pi$ is a Lie algebroid on E whose characteristic foliation lies in $\text{Vert} := \ker p_* \subset TE$.*

In particular, if we restrict the vector bundle $\mathcal{K}$ to a fiber $E_x := p^{-1}(\{x\})$ over any $x \in B$, we obtain a nice Lie algebroid $\mathcal{K}|_{E_x}$ over E_x .

2. EHRESMANN CONNECTIONS

We now introduce a notion of connection [9] for extensions of Lie algebroids.

Definition 2.1. An **Ehresmann connection** on a Lie algebroid extension $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ is a smooth sub-vector bundle $H \subset A_E$ complementary to $\mathcal{K}$ in A_E :

$$\mathcal{K} \oplus H = A_E.$$

We will call H the horizontal distribution of the connection.

Clearly, such a connection always exists and there is an associated notion of **horizontal lifting** $h : \Gamma(A_B) \mapsto \Gamma(H)$: we define $h(\alpha)$ as the unique section of H that π -projects on α .

We might use the terminology of connection in a rather unrestricted way, to refer either to the operator h or to the horizontal sub-bundle H .

Definition 2.2. Given an Ehresmann connection, we define the associated **curvature 2-form** $\omega \in \Omega^2(A_B) \otimes \Gamma(\mathcal{K})$ (otherwise stated, all tensor products are to be taken over the commutative ring $C^\infty(B)$) by the following formula:

$$\omega(\alpha, \beta) := h([\alpha, \beta]_{A_B}) - [h(\alpha), h(\beta)]_{A_E}.$$

The fact that ω takes values in $\Gamma(\mathcal{K})$ is a direct consequence of lemma 1.9. A connection also determines a $C^\infty(B)$ -linear application $\mathcal{D} : \Gamma(A_B) \rightarrow \text{Der}(\mathcal{K})$ defined by:

$$\mathcal{D}_\alpha(\kappa) := [h(\alpha), \kappa]_{A_E}.$$

More precisely for any sections $\alpha, \beta \in \Gamma(A_B)$, any function $g \in C^\infty(B)$ and any sections $\kappa, \kappa_1, \kappa_2 \in \Gamma(\mathcal{K})$:

- (2) $\mathcal{D}_\alpha + \mathcal{D}_\beta = \mathcal{D}_{\alpha+\beta},$
- (3) $\mathcal{D}_{g \cdot \alpha} = g \cdot \mathcal{D}_\alpha,$
- (4) $\mathcal{D}_\alpha([\kappa_1, \kappa_2]_{\mathcal{K}}) = [\mathcal{D}_\alpha(\kappa_1), \kappa_2]_{\mathcal{K}} + [\kappa_1, \mathcal{D}_\alpha(\kappa_2)]_{\mathcal{K}},$
- (5) $\sharp_K(\mathcal{D}_\alpha(\kappa)) = [s_{\mathcal{D}_\alpha}, \kappa]_{TE}.$

Here, the first two equalities express the $C^\infty(B)$ -linearity of $\mathcal{D}$; equality (2) is clear and (3) follows from Proposition 1.10. The last two equations express the fact that $\mathcal{D}$ has values in derivations of $\mathcal{K}$: equation (4) is a consequence of Jacobi identity (iv) and (5) follows from (iii) in Definition 1.1.

As a derivation of $\mathcal{K}$, each $\mathcal{D}_\alpha$ naturally extends to a derivation of the graded algebra $\Gamma(\Lambda^k \mathcal{K})$ of multi-sections of $\mathcal{K}$:

$$\mathcal{D}_\alpha(\kappa_1 \wedge \cdots \wedge \kappa_k) := \sum_{i=1 \dots k} \kappa_1 \wedge \cdots \wedge \mathcal{D}_\alpha(\kappa_i) \wedge \cdots \wedge \kappa_k, \quad (\kappa_i \in \Gamma(\mathcal{K})),$$

and by duality to multi-sections of $\Omega^k(\mathcal{K})$:

$$\langle \mathcal{D}_\alpha \theta, \kappa \rangle := \mathcal{L}_{s_{\mathcal{D}_\alpha}} \langle \theta, \kappa \rangle - \langle \theta, \mathcal{D}_\alpha \kappa \rangle, \quad (\theta \in \Omega^k(\mathcal{K}), \kappa \in \Gamma(\Lambda^k \mathcal{K})).$$

The covariant differential associated to $\mathcal{D}$ will be denoted $\partial_H : \Omega^k(A_B) \otimes \Gamma(\Lambda^l \mathcal{K}) \rightarrow \Omega^{k+1}(A_B) \otimes \Gamma(\Lambda^l \mathcal{K})$ to fit with classical notations (note however that it is related to $\mathcal{D}$ rather than H since two different horizontal distributions can induce the same derivations). It is given by the usual formula:

$$(6) \quad \begin{aligned} \partial_H \theta(\alpha_0 \dots \alpha_k) &:= \sum_{i=0 \dots k} (-1)^i \mathcal{D}_{\alpha_k} \theta(\alpha_0 \dots \hat{\alpha}_i \dots \alpha_k) \\ &+ \sum_{0 \leq i < j \leq k} (-1)^{i+j} \theta([\alpha_i, \alpha_j]_{A_B} \alpha_0 \dots \hat{\alpha}_i \dots \hat{\alpha}_j \dots \alpha_k). \end{aligned}$$

This exact formula can be dualized to define a covariant derivative $\partial_H : \Omega^k(A_B) \otimes \Omega^l(\mathcal{K}) \rightarrow \Omega^{k+1}(A_B) \otimes \Omega^l(\mathcal{K})$ which we will also denote ∂_H .

The curvature $\text{Curv}_{\mathcal{D}}$ of $\mathcal{D}$ is defined by $\text{Curv}_{\mathcal{D}}(\alpha, \beta) := \mathcal{D}_\alpha \circ \mathcal{D}_\beta - \mathcal{D}_\beta \circ \mathcal{D}_\alpha - \mathcal{D}_{[\alpha, \beta]_{A_B}}$. By construction, it is related to the curvature 2-form ω by:

$$(7) \quad \text{Curv}_{\mathcal{D}}(\alpha, \beta)(\kappa) = [\omega(\alpha, \beta), \kappa]_{\mathcal{K}}.$$

Moreover, ω is $\mathcal{D}$ closed in the sense that $\partial_H \omega = 0$, which writes:

$$(8) \quad \oint_{\alpha, \beta, \gamma} \mathcal{D}_\alpha \omega(\beta, \gamma) - \omega([\alpha, \beta]_{A_B}, \gamma), \quad (\alpha, \beta, \gamma \in \Gamma(A_B)).$$

It is a consequence of Jacobi identity (iv).

In order not to confuse $\text{Curv}_{\mathcal{D}}$ with ω we shall refer to ω as the *curvature 2-form*, and to $\text{Curv}_{\mathcal{D}}$ as the *curvature*.

Remark 2.3. The following fact will play an important role in this work: in general, $\mathcal{D}$ might fail to induce an infinitesimal action of A_B on E . However, it is clear in (7) that this failure is still controlled by $\mathcal{K}$. In other words, if we think of $\mathcal{K}$ as a defining a homotopy theory on E , one can see an extension of A_B by $\mathcal{K}$ as an infinitesimal action up to homotopy. Note that this terminology corresponds to a different notion than the one of “representations up to homotopy” appearing for instance in [7].

2.1. Parallel transport. Given a connection, parallel transport along a small A_B -path is well defined, at least locally as we briefly explain now.

Fix a time-dependent section $\alpha^t \in \Gamma(A_B)$. We denote $X_{\mathcal{D}_\alpha}^t$ the linear vector field on $\mathcal{K}$ associated to $\mathcal{D}_{\alpha^t}$ and $\phi_{t,s}^{\mathcal{D}_\alpha}$ its flow. It is *a priori* only defined in a neighborhood of κ in $\mathcal{K}$ and for t close to s .

Now, since the application $\alpha \mapsto \mathcal{D}_\alpha$ is $C^\infty(B)$ -linear, we see that the restriction of $X_{\mathcal{D}_\alpha}^t$ to a fiber $\mathcal{K}_{|E_b}$ over some $b \in B$ only depends on $\alpha^t(b)$. Moreover, by definition $\mathcal{D}_\alpha^t = \text{ad}_{h(\alpha^t)}^{A_E}$ so it has symbol $\sharp_E(h(\alpha^t))$ which we know from Lemma 1.8 to p_* -project onto $\sharp_B(\alpha^t)$. This implies that the restriction of $\phi_{t,0}^{\mathcal{D}_\alpha}$ to $\mathcal{K}_{|E_b}$ only depends on $\alpha^{t'}$ evaluated at the point $\phi_{t',0}^{\sharp_B \alpha}(b)$ for t' in $[0, t]$.

Thus, if $a : I \rightarrow A_B$ is a A_B -path over $\gamma : I \rightarrow B$, one can extend it arbitrarily to a time dependent α^t section of A_B and define $\Phi_{t,0}^a(\kappa) := \phi_{t,0}^{\mathcal{D}_\alpha}(\kappa)$ where $\kappa \in \mathcal{K}_{|E_{\gamma(0)}}$. We know that it is well defined for t small enough (depending on κ) and independent of the choice of α^t .

Definition 2.4. An Ehresmann connection is said to be **complete** if, for any A_B -path a , $\Phi_{t,0}^a(\kappa)$ is well defined for any $\kappa \in \mathcal{K}|_{E_{\gamma(0)}}$ and all $t \in I$.

For a complete connection, we obtain at time $t = 1$ an application:

$$\mathcal{K}|_{E_{\gamma(0)}} \xrightarrow{\Phi_a} \mathcal{K}|_{E_{\gamma(1)}}.$$

called parallel transport or **holonomy** along a . Clearly Φ_a is a Lie algebroid morphism. We shall denote $\phi_a : E_{\gamma(0)} \rightarrow E_{\gamma(1)}$ the base map covered by Φ^a

2.2. Admissible couples. Once fixed a connection, one can decompose any section η of A_E as a sum of elements of the form $\kappa + fh(\alpha)$ with $\kappa \in \Gamma(\mathcal{K})$, $\alpha \in \Gamma(A_B)$ and $f \in C^\infty(E)$ so we get an isomorphism:

$$\Gamma(A_E) = \Gamma(\mathcal{K}) \oplus C^\infty(E) \otimes \Gamma(A_B).$$

The structure of Lie algebroid on A_E is thus entirely determined by the ones on $\mathcal{K}$ and A_B , and by the couple $(\mathcal{D}, \omega)$. Indeed, the brackets are given by:

$$(9) \quad [\kappa_1, \kappa_2]_{A_E} = [\kappa_1, \kappa_2]_{\mathcal{K}},$$

$$(10) \quad [h(\alpha), \kappa]_{A_E} = \mathcal{D}_\alpha \kappa,$$

$$(11) \quad [h(\alpha), h(\beta)]_{A_E} = h([\alpha, \beta]_{A_B}) + \omega(\alpha, \beta),$$

and extended to arbitrary sections using Leibniz rule. The anchor is given by the following formulas:

$$\begin{aligned} \sharp_{A_E}(h(\alpha)) &= s_{\mathcal{D}_\alpha}, \\ \sharp_{A_E}(\kappa) &= \sharp_{\mathcal{K}}(\kappa). \end{aligned}$$

Reciprocally, once fixed a submersion $p : E \rightarrow B$, a Lie algebroid $\mathcal{K}$ over E whose characteristic foliation lies in Vert and a Lie algebroid A_B over B , one shall consider couples $(\mathcal{D}, \omega)$ where:

- $\mathcal{D} : \Gamma(A_B) \rightarrow \text{Der}(\mathcal{K})$ is a $C^\infty(B)$ -linear application such that the symbol $s_{\mathcal{D}_\alpha}$ of $\mathcal{D}_\alpha$ p_* -projects onto $\sharp_B(\alpha)$ for any $\alpha \in \Gamma(A_B)$;
- ω is a 2-form on A_B with values in sections of $\mathcal{K}$: $\omega \in \Omega^2(A_B) \otimes \Gamma(\mathcal{K})$.

Definition 2.5. The couple $(\mathcal{D}, \omega)$ is said to be admissible if $\partial_H \omega = 0$ and $\text{Curv}_{\mathcal{D}} = \text{ad}^{\mathcal{K}} \circ \omega$.

Under the assumption of admissibility, the formulas above define a Lie algebroid structure on $A_E := \mathcal{K} \oplus p^* A_B$ coming with the obvious Ehresmann connection. Note however that given arbitrary Lie algebroids A_B and $\mathcal{K}$, there might not exist an extension of A_B by $\mathcal{K}$.

A natural question is to ask how the choice of the connection influences our construction. Given two connections h and h' , one can form the difference $\Delta := h - h' \in \Omega(A_B) \otimes \Gamma(\mathcal{K})$ and one easily gets the following relations:

$$(12) \quad \mathcal{D}'_\alpha \kappa = \mathcal{D}_\alpha \kappa + [\Delta(\alpha), \kappa]_{\mathcal{K}},$$

$$(13) \quad \omega'(\alpha, \beta) = \omega(\alpha, \beta) + \partial_H \Delta(\alpha, \beta) + [\Delta(\alpha), \Delta(\beta)]_{\mathcal{K}}.$$

where of course $(\mathcal{D}, \omega)$ and $(\mathcal{D}', \omega')$ denote the compatible couples respectively associated to h and h' . Thus one can state the following.

Theorem 2.6. *Let $p : E \rightarrow B$ be a submersion, $\mathcal{K} \rightarrow E$ a Lie algebroid whose characteristic foliation lies in Vert and $A_B \rightarrow B$ a Lie algebroid.*

Extensions of A_B by $\mathcal{K}$ covering p are (up to equivalence) classified by equivalence classes of admissible couples $(\mathcal{D}, \omega)$ under the relation $(\mathcal{D}, \omega) \sim (\mathcal{D}', \omega')$ if there exists $\Delta \in \Omega^1(A_B) \otimes \Gamma(\mathcal{K})$ such that

$$\begin{aligned}\mathcal{D}' &= \mathcal{D} + \text{ad}^{\mathcal{K}} \circ \Delta, \\ \omega' &= \omega + \partial_H \Delta + [\Delta \wedge \Delta]_{\mathcal{K}},\end{aligned}$$

where $[\Delta \wedge \Delta]_{\mathcal{K}} \in \Omega^2(A_B) \otimes \Gamma(\mathcal{K})$ is defined by:

$$[\Delta \wedge \Delta]_{\mathcal{K}}(\alpha, \beta) = [\Delta(\alpha), \Delta(\beta)]_{\mathcal{K}}.$$

Remark 2.7. As for usual considerations concerning extensions, by equivalence of extensions, we mean an algebroid isomorphism $A_E \rightarrow A'_E$ such that the following diagram commutes:

$$\begin{array}{ccccc}\mathcal{K} & \hookrightarrow & A_E & \xrightarrow{\pi} & A_B \\ \downarrow \text{Id}_{\mathcal{K}} & & \downarrow & & \downarrow \text{Id}_{A_B} \\ \mathcal{K} & \hookrightarrow & A'_E & \xrightarrow{\pi'} & A_B.\end{array}$$

In a practical situation, this theorem might be of no real help for classification. However, it gives a good intuition on how Lie algebroid extensions are made.

2.3. Isotropies. For a general extension $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ there is no simple relation between the isotropy Lie algebras $\mathfrak{g}_{p(x)}^B$, $\mathfrak{g}_x^E$ and $\mathfrak{g}_x^{\mathcal{K}}$ for any $x \in E$. In particular, these *do not* fit into an extension of Lie algebras in general, as shows example 2.15. In the next proposition we summarize a few general properties.

Proposition 2.8. *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a Lie algebroid extension. Then the following properties hold:*

- i) $\pi(\mathfrak{g}_x^E) \subset \mathfrak{g}_y^B$;
- ii) $\mathfrak{g}_x^{\mathcal{K}} = \mathfrak{g}_x^E \cap \mathcal{K}_x$;
- iii) $\pi_x^{-1}(\mathfrak{g}_y^B) = (\sharp_E)_x^{-1}(\text{Vert}_x)$;

for any $x \in E$, $y = p(x)$.

Definition 2.9. We will say that an extension is **clean** provided the restriction of π to $\mathfrak{g}_x^E$ surjects onto $\mathfrak{g}_{p(x)}^B$ for any $x \in E$.

As easily seen, in the case of a clean extension the restriction of π to $\mathfrak{g}_x^E$ induces an extension of Lie algebras:

$$\mathfrak{g}_x^{\mathcal{K}} \hookrightarrow \mathfrak{g}_x^E \twoheadrightarrow \mathfrak{g}_{p(x)}^B.$$

Moreover, clean extensions enjoy the following property: each orbit $\mathcal{O}_E$ of A_E fibers over an orbit $\mathcal{O}_B$ of A_B , the typical fiber having as connected components orbits of $\mathcal{K}$ (this follows from (i) and (iii) in Proposition 2.8).

Once fixed an orbit $\mathcal{O}_E$ of A_E , one can always choose a Ehresmann connection such that $h(\mathfrak{g}_y^B) \subset \mathfrak{g}_x^E$ for any $y \in B$, $x \in p^{-1}(y) \cap \mathcal{O}_E$. Note however that this might not be possible for all $e \in E$ in general.

For such a connection, the restriction of the curvature 2-form ω to sections of the isotropy Lie algebra bundle $\mathfrak{g}_{|\mathcal{O}_B}^B$ takes values in $\mathfrak{g}_{|\mathcal{O}_E}^{\mathcal{K}}$.

2.4. Semi-direct products.

Definition 2.10. An Lie algebroid extension will be said of **semi-direct product type** provided it admits a connection whose curvature 2-form vanishes: $\omega = 0$. In particular $\mathcal{D}$ has no curvature as well by (7). Once fixed such a connection, we shall write $A_E = A_B \ltimes \mathcal{K}$.

Since the curvature of $\mathcal{D}$ vanishes, the holonomy along an A_B -path depends only on its A_B -homotopy class. This follows from the fact that $\mathcal{D} : \Gamma(A_B) \rightarrow \text{Der}(\mathcal{K})$ is a Lie algebra morphism and can be proved using arguments similar to those in the proof of proposition A.1 (see also equation (36)).

2.5. Abelian Extensions.

Definition 2.11. An **abelian extension** is an extension whose kernel $\mathcal{K}$ is a bundle of abelian Lie algebras over E .

If one fixes a connection, because of condition (7), we have an action of A_B on $E \rightarrow B$, plus a representation of the associated action algebroid $A_B \ltimes E$ on $\mathcal{K}$. Note however that $A_B \ltimes \mathcal{K}$ and A_E may differ in general since the curvature 2-form ω might not vanish. In practice, this means that one gets the bracket on $A_B \ltimes \mathcal{K}$ by the one on A_E by simply deleting the term involving ω in (11).

In the case $E = B$, then $\mathcal{K}$ is a representation of A_B ; the curvature 2-form is submitted to the only condition (8) which exactly means that it is a closed form on A_B with values in the representation $\mathcal{K}$. In the Theorem 2.6 the bracket $[\cdot, \cdot]_{\mathcal{K}}$ vanishes, so it is easily seen that only the cohomology class of ω determines the extension up to equivalence. In particular, it vanishes if and only if the extension is of semi-direct product type.

Definition 2.12. A **central extension** is an abelian extension with $E = B$, and $\mathcal{K}$ the trivial bundle of abelian Lie algebras over B , $\mathcal{K} = \mathbb{R}^n \times B$ on which A_B acts trivially.

To be more precise, the condition that A_B acts trivially means that $\mathcal{D}_\alpha \kappa = \mathcal{L}_{\sharp_B(\alpha)} \kappa$ where we see a section of $\mathcal{K}$ as an application $\kappa : B \rightarrow \mathbb{R}^n$.

The only condition on the curvature form ω in order to define an extension is to be d_A -closed. Therefore central extensions on A_B are in one to one correspondence with closed 2-form with values in $\mathbb{R}^n$.

We shall refer to [8] for a nice account of such constructions in the context of Poisson and Jacobi manifolds.

2.6. Examples.

Example 2.13. *Extensions of Lie algebras.* When both E and B are reduced to a point, $E = B = \{*\}$, both A_E and A_B are Lie algebras, and we have an extension of Lie algebras. In that case, the notions of abelian and central extensions coincide with the usual ones.

Example 2.14. *Submersions.* A submersion $p : E \rightarrow B$ realizes TE as an extension of TB by the vertical bundle: $\text{Vert} \hookrightarrow TE \twoheadrightarrow TB$.

Example 2.15. Infinitesimal Actions. Consider an infinitesimal action of a Lie algebra $\mathfrak{g}$ on a smooth manifold M that is a Lie algebra morphism $\mathfrak{g} \rightarrow \mathfrak{X}(M)$; we denote $\mathfrak{g} \ltimes M$ the corresponding action algebroid. Then one gets an extension by taking $E := M$, $A_E := \mathfrak{g} \ltimes M$ and $A_B = \mathfrak{g}$ seen as an algebroid over a point $B = \{*\}$. In that case, the kernel is trivial $\mathcal{K} = E \times \{0\}$ and there is a canonically defined Ehresmann connection. In fact one can *define* an infinitesimal action as an extension of a Lie algebra with trivial kernel.

More generally, given an algebroid $A_B \rightarrow B$ and an action of A_B on a fibration $E \rightarrow B$, the associated action algebroid $A_B \ltimes E \rightarrow E$ is an extension of A_B with vanishing kernel: $\mathcal{K} = E \times \{0\}$. Reciprocally, an extension of an algebroid A_B with vanishing kernel $\mathcal{K}$ is naturally associated to an action of A_B on E .

In this situation, one can also form the semi-direct product $A_B \ltimes \text{Vert} \rightarrow E$, which is an extension of A_B by $\text{Vert} := \ker p_* \subset TE$. See [14] for more details.

Example 2.16. Pull-back Lie algebroids. Given a submersion $E \rightarrow B$ and an algebroid $A_B \rightarrow B$ over B , the pull-back algebroid $p^!A_B$ is defined by:

$$\pi^!A_B := \{(X, \alpha) \in TE \times A_B / p_*X = \sharp_B \alpha\}.$$

Clearly, the projection on the second factor $\pi : p^!A_B \rightarrow A_B$ is a surjective algebroid morphism with kernel $\mathcal{K} = \text{Vert}$, where $\text{Vert} \subset TE$ denotes the vertical bundle: $\text{Vert} := \ker p_*$. A connection amounts to a partially defined connection on $\text{Im } \sharp_B$, one can for instance choose a usual Ehresmann connection on $E \rightarrow B$ and compose it with $\sharp_B$.

Example 2.17. Fibered Lie algebroids. In the case of an extension $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ where $A_B = \{0\} \times B$, A_E coincides with $\mathcal{K}$ which stands as a fibered Lie algebroid over $p : E \rightarrow B$.

Example 2.18. Extension by bundles of Lie algebras. In the case of an extension $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ where E and B do coincide: $E = B$, it is easily seen that $\mathcal{K}$ has vanishing anchor so it is a bundle of Lie algebras.

Example 2.19. The Atiyah exact sequence. A transitive Lie algebroid $A \rightarrow B$ naturally fits into an extension of $A_B = TB$. Here, $B = E$ and the projection π is given by the anchor $\pi = \sharp$. Then $\mathcal{K} := \ker \sharp$ is a bundle of Lie algebras. An Ehresmann connection is then just a section of the anchor, and the corresponding curvature 2-form is an element $\omega \in \Omega^2(B) \otimes \Gamma(\ker \sharp)$.

A regular Lie algebroid $A \rightarrow B$ with foliation $F \subset TB$ can be treated similarly as an extension of F by a bundle of Lie algebras.

Example 2.20. Neighborhood of an orbit. The following is a straightforward adaptation of the construction in [18].

Consider an algebroid $A_E \rightarrow E$, and let B denote one of its orbits. Assume B admits a tubular neighborhood so that, up to a shrinking of E , one is given a smooth submersion $p : E \rightarrow B$, as usual we denote $\text{Vert} := \ker p_*$.

Since along B we have $T_BE = \text{Vert}|_B \oplus \text{Im } \sharp_E$, the following still holds in a whole neighborhood of B : $TE = \text{Vert} + \text{Im } \sharp_E$. Thus, shrinking E if necessary, one obtains one obtains a surjective bundle map $A_E \rightarrow TB$ by compositing p_* with $\sharp_E$. It is clearly an algebroid morphism since both p_*

and $\sharp_E$ are. In that case, $\mathcal{K}$ is just the transverse Lie algebroid for each fiber of the projection.

Example 2.21. *Extensions of tangent bundles* Here is a rather general procedure to obtain an extension of a tangent bundle $A_B = TB$, directly inspired by the so-called Yang-Mills-Higgs settings for coupling Dirac structures, see for instance [19], [1] and references therein.

Consider a G -principal bundle $P \rightarrow B$, and an inner action of a group G on an algebroid $A_F \rightarrow F$. By definition, this means that we are given an action $\rho : G \rightarrow \text{Aut}(A_F)$ of G on A_F by Lie algebroid automorphisms, and a G -equivariant algebroid morphism:

$$\psi : \mathfrak{g} \ltimes F \rightarrow A_F,$$

such that the derivation $\text{ad}_{\psi(\xi)}^{A_F}$ has infinitesimal generator $\frac{d}{dt}\rho_{\exp t\xi}$.

Then we can build an extension of TB as follows: fix a principal connection θ on P , then we can use parallel transport to obtain a derivation of the fibered Lie algebroid $\mathcal{K} := P \times_G A_F$:

$$\mathcal{D} : \mathfrak{X}(B) \rightarrow \text{Der}(\mathcal{K}).$$

Define now $\omega \in \Omega^2(B) \otimes \Gamma(\mathcal{K})$ by the following formula:

$$\omega(X, Y) := (P \times_G \psi)(\text{Curv}_\theta(X, Y)),$$

where $\text{Curv}_\theta : \Omega^2(B) \otimes \Gamma(P \times_G \mathfrak{g})$ denotes the curvature of θ as a principal connection, and $P \times_G \psi$ denotes the fibered map:

$$P \times_G \psi : P \times_G \mathfrak{g} \rightarrow P \times_G A_F = \mathcal{K}.$$

It is easily checked that $(\mathcal{D}, \omega)$ induces an extension of TB by $\mathcal{K}$ whose isomorphism class is independent of the choice of θ .

In such a situation, it is possible to work out “by hands” a concrete description of the integrating groupoid, involving only $\mathcal{G}(A_F)$ and bundles associated to P . This will be explained in a separate work [2]

3. COHOMOLOGY

In this section, we describe a spectral sequence converging to the cohomology of an extension. By choosing a connection, we obtain formulas for the successive coboundary operators, which we believe makes it easier to understand. A short example is then discussed.

3.1. The spectral sequence. Given an extension $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$, there is a natural filtration of the complex of cochains $C^k := \Omega^k(A_E)$:

$$C^k =: F_0 C^k \supset \dots \supset F_q C^k \supset F_{q+1} C^k \supset \dots \supset F_{k+1} C^k := \{0\},$$

where the generic term is defined as follows:

$$F_q C^k := \{\alpha \in \Omega^k(A_E), i_v \alpha = 0 \ \forall v \in \Gamma(\Lambda^{k-q+1} \mathcal{K})\}.$$

Since this filtration is bounded, one gets a spectral sequence converging to $H^\bullet(A_E)$ by the usual construction. The only thing one actually has to check is that $d_{A_E}(F_q C^k) \subset F_q C^{k+1}$ which we will leave to the reader (it also can be seen as a consequence of next proposition).

Remark 3.1. We emphasize the change of notations here: up to now, we used Greek letters to denote sections of the different algebroids and Latin ones for vector fields. In this section, Greek letters will be used to denote k -forms on Lie algebroids while latine capitals will be used to refer to multi-section of algebroids, as we think of them as generalized multi-vector fields.

Let us choose an Ehresmann connection (fixed once for all) and recall the identification $\Gamma(A_E) = C^\infty(E) \otimes \Gamma(A_B) \oplus \Gamma(\mathcal{K})$ induced by the splitting $A_E = \mathcal{K} \oplus H$. Dually one has $\Omega(A_E) = \Omega(A_B) \otimes C^\infty(E) \oplus \Omega(\mathcal{K})$. Extending this principle to multi-forms on A_E , one gets an isomorphism:

$$(14) \quad \Omega^k(A_E) = \bigoplus_{p+q=k} \Omega^p(A_B) \otimes \Omega^q(\mathcal{K}),$$

where the projection $\theta^{p,q}$ of some $\Theta \in \Omega^k(A_E)$ on $\Omega^p(A_B) \otimes \Omega^q(\mathcal{K})$ is given by the following formula:

$$(15) \quad \langle \theta^{p,q}(v_1 \dots v_p), w_1 \dots w_q \rangle = \langle \Theta, h(v_1) \dots h(v_p), w_1 \dots w_q \rangle,$$

where $v_i \in \gamma(A_B), w_i \in \Gamma(\mathcal{K})$. The reason why this identification is particularly suiting for our purposes is that the filtration $F_q C^k$ comes as successive truncations:

$$F_q C^k = \bigoplus_{q \leq p \leq k} \Omega^p(A_B) \otimes \Omega^{k-p}(\mathcal{K}).$$

In particular we have $E_0^{p,q} := F_q C^{p+q} / F_{q+1} C^{p+q} = \Omega^p(A_B) \otimes \Omega^q(\mathcal{K})$. In next proposition, we decompose the co-differential operator d_{A_E} under the identification (14), this gives explicit formulas for the successive coboundary operators involved in the spectral sequence.

Proposition 3.2. *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be an extension and $A_E = \mathcal{K} \oplus H$ a fixed Ehresmann connection. Then, under the identification (14) the coboundary operator d_{A_E} on $\Omega^\bullet(A_E^*)$ decomposes as a sum $d_{A_E} = \delta_{0,1} + \delta_{1,0} + \delta_{2,-1}$ where $\delta_{i,j}$ have bi-degree (i,j) . The restrictions of these operators to $\Omega^p(A_B) \otimes \Omega^q(\mathcal{K})$ are given by the following formulas:*

$$(\delta_{0,1}\theta)(v_1 \dots v_p) = (-1)^p d_{\mathcal{K}}(\theta(v_1 \dots v_p)),$$

which is a coboundary operator naturally extending $d_{\mathcal{K}}$ to $\Omega^p(A_B) \otimes \Omega^q(\mathcal{K})$;

$$(16) \quad (\delta_{1,0}\theta)(v_0 \dots v_p) = \sum_{i=0 \dots p} (-1)^i \mathcal{D}_{v_i}(\theta(v_1 \dots \widehat{v}_i \dots v_p)) \\ + \sum_{0 \leq i < j \leq n} (-1)^{i+j} \theta([v_i, v_j]_{A_B}, v_1 \dots \widehat{v}_i \dots \widehat{v}_j \dots v_p),$$

which is the covariant derivative ∂_H associated to the Ehresmann connection as described in Section 2;

$$(\delta_{2,-1}\theta)(v_0 \dots v_{p+1}) = (-1)^{p+1} \sum_{0 \leq i < j \leq p+1} (-1)^{i+j} i_{\omega(v_i, v_j)} \theta(v_0 \dots \widehat{v}_i \dots \widehat{v}_j \dots v_{p+1}),$$

where $\omega \in \Omega^2(A_B) \otimes \Gamma(\mathcal{K})$ is the curvature 2-form of the connection.

Proof. Let us fix $\theta^{p,q} \in \Omega^p(A_B) \otimes \Omega^q(\mathcal{K})$ and denote $\Theta \in \Omega^{p+q}(A_E)$ the corresponding $p+q$ -form on A_E through the isomorphism (14). By construction,

Θ has the following property: for any $v \in \Gamma(\Lambda^s A_B)$ and $v \in \Gamma(\Lambda^t A_B)$ with $s + t = p + q$, the evaluation $\langle \Theta, h(v) \wedge v \rangle$ vanishes whenever $s > p$ or $t > q$.

The different components in $\Omega^s(A_B) \otimes \Omega^t(\mathcal{K})$, where $s + t = p + q + 1$, of $d_{A_E} \Theta$ are obtained using (15) so we just have to compute $\langle d_{A_E} \Theta, v \wedge w \rangle$ for any $v \in \Gamma(\Lambda^s A_B)$, $w \in \Gamma(\Lambda^t \mathcal{K})$, then relate it to $\theta^{p,q}$.

Let us first express $\delta_{0,1}$. We choose sections $w_0 \dots w_q$ of $\mathcal{K}$ and $v_1 \dots v_p$ of A_B and compute:

$$\begin{aligned}
& \langle d_{A_E} \Theta, h(v_1) \dots h(v_p), w_0 \dots w_q \rangle \\
&= \sum_{i=1 \dots p} (-1)^{i+1} \mathcal{L}_{\#_E h(v_i)} \langle \Theta, h(v_1) \dots \widehat{h(v_i)} \dots h(v_p), w_0 \dots w_q \rangle \\
&+ \sum_{1 \leq i < j \leq p} (-1)^{i+j} \langle \Theta, [h(v_i), h(v_j)]_{A_E}, h(v_1) \dots \widehat{h(v_i)} \dots \widehat{h(v_j)} \dots h(v_p), w_0 \dots w_q \rangle \\
&\quad + (-1)^p \sum_{i=0 \dots q} (-1)^i \mathcal{L}_{\#_E w_i} \langle \Theta, h(v_1) \dots h(v_p), w_0 \dots \widehat{w_i} \dots w_q \rangle \\
&\quad + \sum_{0 \leq i < j \leq q} (-1)^{i+j} \langle \Theta, [w_i, w_j]_{A_E}, h(v_1) \dots h(v_p), w_0 \dots \widehat{w_i} \dots \widehat{w_j} \dots w_q \rangle \\
&+ (-1)^{p+1} \sum_{\substack{i=1 \dots p \\ j=0 \dots q}} (-1)^{i+j} \langle \Theta, [h(v_i), w_j]_{A_E}, h(v_1) \dots \widehat{h(v_i)} \dots h(v_p), w_0 \dots \widehat{w_j} \dots w_q \rangle.
\end{aligned}$$

Here, the two first terms vanish since Θ is evaluated on $q + 1$ sections of $\mathcal{K}$, as well as the last term according to lemma 1.9. In the two remaining terms, it is recognized the coboundary operator for the Lie algebroid $\mathcal{K}$ and one obtains the desired expression for $\delta_{0,1}$. Note that one can also write $\delta_{0,1}$ as $\delta_{0,1} = \sum (-1)^p \text{id} \otimes d_{\mathcal{K}}$.

We now compute $\delta_{1,0}$. For any sections $v_0 \dots v_p$ of A_B and $w_1 \dots w_q$ of $\mathcal{K}$, we have:

$$\begin{aligned}
& \langle d_{A_E} \Theta, h(v_0) \dots h(v_p), w_1 \dots w_q \rangle \\
&= \sum_{i=0 \dots p} (-1)^i \mathcal{L}_{\#_E h(v_i)} \langle \Theta, h(v_0) \dots \widehat{h(v_i)} \dots h(v_p), w_1 \dots w_q \rangle \\
&+ \sum_{0 \leq i < j \leq p} (-1)^{i+j} \langle \Theta, [h(v_i), h(v_j)]_{A_E}, h(v_0) \dots \widehat{h(v_i)} \dots \widehat{h(v_j)} \dots h(v_p), w_1 \dots w_q \rangle \\
&\quad + (-1)^p \sum_{i=1 \dots q} (-1)^i \mathcal{L}_{\#_E w_i} \langle \Theta, h(v_0) \dots h(v_p), w_1 \dots \widehat{w_i} \dots w_q \rangle \\
&\quad + \sum_{1 \leq i < j \dots \leq q} (-1)^{i+j} \langle \Theta, [w_i, w_j]_{A_E}, h(v_0) \dots h(v_p), w_1 \dots \widehat{w_i} \dots \widehat{w_j} \dots w_q \rangle \\
&+ (-1)^p \sum_{\substack{i=0 \dots p \\ j=1 \dots q}} (-1)^{i+j} \langle \Theta, [h(v_i), w_j]_{A_E}, h(v_0) \dots \widehat{h(v_i)} \dots h(v_p), w_1 \dots \widehat{w_j} \dots w_q \rangle.
\end{aligned}$$

In this sum, the third and fourth terms vanish since Θ is evaluated on $p+1$ sections of H . Then, taking apart the first and last terms, one gets:

$$\begin{aligned}
& \sum_{i=0 \dots p} (-1)^i \mathcal{L}_{\sharp_E h(v_i)} \langle \Theta, h(v_0) \dots \widehat{h(v_i)} \dots h(v_p), w_1 \dots w_q \rangle \\
& + (-1)^p \sum_{\substack{i=0 \dots p \\ j=1 \dots q}} (-1)^{i+j} \langle \Theta, \mathcal{D}_{v_i} w_j, h(v_0) \dots \widehat{h(v_i)} \dots h(v_p), w_1 \dots \widehat{w_j} \dots w_q \rangle \\
& = \sum_{i=0 \dots p} (-1)^i \mathcal{L}_{\sharp_E h(v_i)} \langle \theta^{p,q}(v_0 \dots \widehat{v_i} \dots v_p), w_1 \dots w_q \rangle \\
& + \sum_{\substack{i=0 \dots p \\ j=1 \dots q}} (-1)^{i+j} \langle \theta^{p,q}(v_0 \dots \widehat{v_i} \dots v_p), \mathcal{D}_{v_i} w_j, w_1 \dots \widehat{w_j} \dots w_q \rangle \\
& = \sum_{i=0 \dots p} (-1)^i \langle \mathcal{D}_{v_i} \theta^{p,q}(v_0 \dots \widehat{v_i} \dots v_p), w_1 \dots w_q \rangle.
\end{aligned}$$

Considering now the second term, we know that the following expression:

$$\langle \Theta, \omega(v_i, v_j), h(v_0) \dots \widehat{h(v_i)} \dots \widehat{h(v_j)} \dots h(v_p), w_1 \dots w_q \rangle$$

vanishes because Θ is evaluated on $q+1$ section of $\mathcal{K}$. Since ω precisely measures the difference $\omega(v_i, v_j) = [h(v_i), h(v_j)]_{A_E} - h([v_i, v_j]_{A_B})$ one deduces that

$$\begin{aligned}
& \sum_{0 \leq i < j \leq p} (-1)^{i+j} \langle \Theta, [h(v_i), h(v_j)]_{A_E}, v_0 \dots \widehat{v_i} \dots \widehat{v_j} \dots v_p, w_1 \dots w_q \rangle \\
& = \sum_{0 \leq i < j \leq p} (-1)^{i+j} \langle \Theta, h([v_i, v_j]_{A_B}), v_0 \dots \widehat{v_i} \dots \widehat{v_j} \dots v_p, w_1 \dots w_q \rangle \\
& = \sum_{0 \leq i < j \leq p} (-1)^{i+j} \langle \theta^{p,q}([v_i, v_j]_{A_B}), v_0 \dots \widehat{v_i} \dots \widehat{v_j} \dots v_p, w_1 \dots w_q \rangle.
\end{aligned}$$

So we obtain the announced formula for $\delta_{1,0}$. It appears to coincide with the covariant derivative ∂_H associated to the connection.

Let us now express $\delta_{2,-1}$. For any sections $w_1 \dots w_{q-1}$ of $\mathcal{K}$ and $v_0 \dots v_{p+1}$ of A_B we check that:

$$\begin{aligned}
& \langle d_{A_E} \Theta, h(v_0) \dots h(v_{p+1}), w_1 \dots w_{q-1} \rangle \\
& = \sum_{i=0 \dots p+1} (-1)^i \mathcal{L}_{\sharp_E h(v_i)} \langle \Theta, h(v_0) \dots \widehat{h(v_i)} \dots h(v_{p+1}), w_1 \dots w_{q-1} \rangle \\
& + \sum_{0 \leq i < j \leq p+1} (-1)^{i+j} \langle \Theta, [h(v_i), h(v_j)]_{A_E}, h(v_0) \dots \widehat{h(v_i)} \dots \widehat{h(v_j)} \dots h(v_p), w_1 \dots w_{q-1} \rangle \\
& + (-1)^{p+1} \sum_{j=1 \dots q} (-1)^j \mathcal{L}_{\sharp_E w_j} \langle \Theta, h(v_0) \dots h(v_{p+1}), w_1 \dots \widehat{w_j} \dots w_{q+1} \rangle \\
& + \sum_{1 \leq i < j \leq q-1} (-1)^{i+j} \langle \Theta, [w_i, w_j]_{A_E}, h(v_0) \dots h(v_{p+1}), w_1 \dots \widehat{w_i} \dots \widehat{w_j} \dots w_{q-1} \rangle \\
& + (-1)^{p+1} \sum_{\substack{i=0 \dots p+1 \\ j=1 \dots q-1}} (-1)^{i+j} \langle \Theta, [h(v_i), w_j]_{A_E}, h(v_0) \dots \widehat{h(v_i)} \dots h(v_{p+1}), w_1 \dots \widehat{w_j} \dots w_{q-1} \rangle.
\end{aligned}$$

Here, it is easily seen that only the second term does not vanish. To deal with it, one notices that the expression

$$\langle \Theta, h([v_i, v_j]_{A_B}), h(v_0) \dots \widehat{h(v_i)} \dots \widehat{h(v_j)} \dots h(v_p), w_1 \dots w_{q-1} \rangle$$

vanishes since Θ is evaluated on $p+1$ section of H , and since ω precisely measures the difference $\omega(v_i, v_j) = [h(v_i), h(v_j)]_{A_E} - h([v_i, v_j]_{A_B})$ one gets the expression for $\delta_{2,-1}$ stated in the proposition.

To conclude one has to check that $d_A \Theta$ has no other components than those three mentioned in this proof, but this should be clear by what has been done so far. $\square$

In order to keep notations meaningful, in the sequel, we will write respectively ∂_K , ∂_H , and ∂_ω rather than $\delta_{0,1}$, $\delta_{1,0}$ and $\delta_{2,-1}$. Using the bi-graduation and the fact that $d_{A_E} \circ d_{A_E} = 0$ one easily gets the following useful formulas.

Proposition 3.3. *Under the hypothesis of proposition 3.2 and notations above, the following relations holds:*

- i) $\partial_K \circ \partial_K = 0,$
- ii) $\partial_K \circ \partial_H + \partial_H \circ \partial_K = 0,$
- iii) $\partial_K \circ \partial_\omega + \partial_H \circ \partial_H + \partial_\omega \circ \partial_K = 0,$
- iv) $\partial_H \circ \partial_\omega + \partial_\omega \circ \partial_H = 0,$
- v) $\partial_\omega \circ \partial_\omega = 0.$

3.2. Interpretation. Let us now explain how these results can give some sort of geometric approach to this cohomology. Here, we will carry out the usual construction of a spectral sequence associated to a filtration, using standard notations.

We explained already why:

$$E_0^{p,q} := F_q^{p,q} \Omega(A_E)^{p+q} / F_{q+1}^{p,q} \Omega^{p+q}(A_E) = \Omega^p(A_B) \otimes \Omega^q(K),$$

and we have $d_0 = \text{id} \otimes d_K$ according to the Proposition 3.2.

Now the first sheet $E_1^{p,q}$ of the spectral sequence can be thought of as forms on A_B with values in the cohomology $H^*(K)$ of K because:

$$E_1^{p,q} := \Omega^p(A_B) \otimes Z^q(K) / \Omega^p(A_B) \otimes B^q(K),$$

where we denoted:

$$Z^q(K) := \{\mu \in \Omega^q(K) : d_K \mu = 0\},$$

$$B^q(K) := \{\mu \in \Omega^q(K) : \mu = d_K \nu\}.$$

The coboundary operator d_1 on $E_1^{p,q}$ is the quotient of the map ∂_H . Let us see why $d_1 \circ d_1 = 0$: an element in $E_1^{p,q}$ is represented by some $\theta \in \Omega^p(B) \otimes Z^q(K)$ thus, using (iii) in proposition 3.3, one has:

$$-\partial_H \circ \partial_H \theta = \partial_K \circ \partial_\omega \theta + \partial_\omega \circ \partial_K \theta = \partial_K \circ \partial_\omega \theta,$$

which indeed represents a trivial element in $E_1^{p+1,q}$.

Now, in order to understand the second sheet $E_2^{p,q}$, one shall think of elements in $H^*(K)$ as sections of a vector bundle over B . This makes sense since $H^*(K)$ carries a natural structure of $C^\infty(B)$ -module (this because d_K is $C^\infty(B)$ -linear). From this point of view ∂_H induces a representation of A_B on $H^*(K)$ by setting $D_X([\theta]_K) := [\partial_H \theta(X)]_K$. The equality (ii) in proposition 3.3 ensures that D is well defined, and the last computation

with $p = 0$ that it is indeed a representation. Thus the second sheet $E_2^{p,q}$ is as close as one can get to a cohomology of A_B with values in the representation $H^*(\mathcal{K})$.

The last sheet $E_3^{p,q}$ specifically involves ∂_ω , it will be the crucial step distinguishing the cohomologies of two extensions with same derivative $\mathcal{D}$ but distinct curvatures forms ω, ω' .

We stress the fact that the spaces we are dealing with here can be very singular. In some examples though, this construction allows to carry the computation of the cohomology through. Of course, this assumes a good knowledge of the cohomologies of A_B and $\mathcal{K}$.

Example 3.4. Applying these results to an action Lie algebroid (see Example 2.15), we get the well-known fact that the cohomology of $g \ltimes M$ is the same as the Chevalley Eilenberg cohomology of $\mathfrak{g}$ with values in the $\mathfrak{g}$ -module $C^\infty(M)$.

Example 3.5. In the case of a fibration $E \rightarrow B$ (see Example 2.14), one obtains the usual Leray Serre spectral sequence, involving vertical de Rham cohomology on E and the de Rham cohomology of the base.

Example 3.6. In the case of Poisson cohomology relative to a symplectic leaf, which is a particular case of cohomology of a transitive Lie algebroid, one obtains the same formula as in [11].

3.3. An example of computation. Let us present a simple but non-trivial example where the computation of the cohomology can be carried out.

We consider a Poisson structure on $E := \mathbb{R}^+ \times S^2$ whose symplectic leaves are $\{r\} \times S^2$ endowed with the symplectic form $f\omega_0$, where f is a nowhere vanishing function of $r \in \mathbb{R}^+$ and ω_0 the usual symplectic form on S^2 . Recall that these settings allow the simplest examples of non-integrable Lie algebroids. The associated Poisson cohomology of E was computed in [10], we shall also mention [21] for earlier work on regular Poisson structures. We present here an approach using extensions.

We now set $A_E = T^*E = T^*\mathbb{R}^+ \times T^*S^2$, and we let $p : \mathbb{R}^+ \times S^2 \rightarrow S^2$ be projection on the second factor and $\pi : T^*E \rightarrow T^*S^2$ be the composition of $\sharp_E$ with p_* . Thus T^*E is an extension of $A_B = T^*S^2$ by $\mathcal{K} = T^*\mathbb{R}^+ \times S^2$; note that forms on $\mathcal{K}$ are p -vertical multi-vector fields.

One defines an Ehresmann connection by letting: $h(X) := fi_X\omega_0$. Thus we see that the identification $T^*E = T^*\mathbb{R}^+ \times T^*S^2$ also corresponds with the decomposition $A_E = \mathcal{K} \oplus H$. The curvature form $\omega \in \Omega^2(B) \otimes \Gamma(\mathcal{K})$ is easily computed:

$$\omega(X, Y) = \omega_0(X, Y) \otimes f' dr.$$

Let us now compute the associated cohomology. The kernel $\mathcal{K}$ is an abelian algebroid (with trivial anchor and bracket) so that $d_{\mathcal{K}} = 0$. This implies $E_0^{p,q} = E_1^{p,q} = \Omega^p(TB) \otimes \Omega^q(\mathcal{K})$.

We now compute the coboundary operator d_1 on $E_1^{0,0} = C^\infty(E)$ and $E_1^{0,1} = \Omega^1(\mathcal{K})$. Using formulas in proposition 3.2 we get:

$$\begin{aligned} (\partial_H g)(X) &= \mathcal{L}_X g, \\ (\partial_H g \cdot \partial r)(X) &= (\mathcal{L}_X g) \partial r, \end{aligned}$$

for any $g \in C^\infty(E)$. So we see that $\partial_H(g) = 0$ or $\partial_H(g\partial r) = 0$ if and only if g only depends on r . It follows that $E_2^{0,0}$ and $E_2^{0,1}$ can be identified to $C^\infty(\mathbb{R}^+)$.

The explicit formula for d_1 on $E_1^{1,0} = \Omega^1(TB) \otimes C^\infty(E)$ is given by:

$$(\partial_H \alpha)(X, Y) = \mathcal{L}_X \alpha(Y) - \mathcal{L}_Y \alpha(X) - \alpha([X, Y]), \quad (\alpha \in E_1^{1,0}).$$

Here $\alpha(X)$, $\alpha(Y)$, and $\alpha([X, Y])$ are function on $\mathbb{R}^+ \times S^2$ but since an element α of $E_1^{1,0}$ is the same as a smooth family of 1-forms on S^2 with parameter r , we see that $\partial_H \alpha = 0$ if and only if it determines a family of closed forms. It can be made into a family of exact 1-forms because $H^1(S^2) = \{0\}$ and this can be done in a smooth way with respect to r ; so we conclude that $E_2^{1,0} = \{0\}$. For the same reason, if we use the pairing with dr to identify $E_1^{1,1}$ with $\Omega^2(B) \otimes C^\infty(E)$, we one get the same formula for ∂_H as above, thus $E_1^{1,1} = \{0\}$ as well.

The same kind of reasoning applies to compute $E_2^{2,0}$: we know that integration of 2-forms over S^2 induces an isomorphism $H^2(S^2) \rightarrow \mathbb{R}$. A ∂_H -closed element in $E_1^{2,0} = \Omega^2(B) \otimes C^\infty(E)$ being the same thing as a family of closed 2-forms on S^2 with parameters in $\mathbb{R}^+$, one sees that integration over S^2 with fixed parameter yields an isomorphism $E_2^{2,0} \simeq C^\infty(\mathbb{R}^+)$ given by:

$$[g \cdot \omega_0] \mapsto (r \mapsto \int_{S^2} g(r, \cdot) \omega_0).$$

For the same reasons, one gets has isomorphism $E_2^{2,1} \simeq C^\infty(\mathbb{R}^+)$ via

$$[\omega_0 \otimes g\partial r] \mapsto (r \mapsto \int_{S^2} g(r, \cdot) \omega_0).$$

These identifications are convenient for our purposes because, as easily checked, the explicit formula for $d_2 := [d_\omega] : E_2^{0,1} \rightarrow E_2^{2,0}$ is just multiplication by f' . Since d_2 vanishes on $E_2^{p,q}$ for $(p, q) \neq (2, -1)$ the computation is easily concluded. We finally obtain the Poisson cohomology of T^*E as follows:

$$\begin{aligned} H^0(T^*E) &= C^\infty(\mathbb{R}^+), \\ H^1(T^*E) &= \{g \in C^\infty(\mathbb{R}^+), f'g = 0\}, \\ H^2(T^*E) &= C^\infty(\mathbb{R}^+)/f' \cdot C^\infty(\mathbb{R}^+), \\ H^3(T^*E) &= C^\infty(\mathbb{R}^+). \end{aligned}$$

4. INTEGRATION

We will here take advantage of a connection in order to describe A_E -paths and A_E -homotopies, obtaining an alternative description of $\mathcal{G}(A_E)$. From now on, the connection will always be assumed to be **complete**.

4.1. Splitting A_E -paths and A_E -homotopies. Since the essential feature of a Ehresmann connection is to split A_E into a direct sum

$$A_E = \mathcal{K} \oplus H,$$

it is natural, in order to describe $\mathcal{G}(A_E)$, to try to decompose A_E -paths accordingly. However, for a A_E -path $a : I \rightarrow A_E$ covering $\gamma : I \rightarrow E$, the corresponding decomposition is the following:

$$a(t) = a_V(t) + h(a_B(t)),$$

where $a_B := \pi \circ a$ and $a_V : I \rightarrow \mathcal{K}$. This is not satisfying because none of the two terms above is an A_E -path.

Thus, it is better to consider the couple $(\dot{\gamma}_B, a_K)$ where

$$(17) \quad a_K(t) := \Phi_{0,t}^{\gamma_B} \circ a_V(t).$$

Here $\Phi_{t,0}^{\gamma_B} : \mathcal{K}_{|E_{\gamma_B(0)}} \mapsto \mathcal{K}_{|E_{\gamma_B(t)}}$ denotes the parallel transport along a_B . Indeed, it is easily checked that a_K is a A_E -path over $\gamma_K(t) = \phi_{0,t}^{\gamma_B} \circ \gamma(t)$ lying in $E_{\gamma_B(0)}$. Moreover the correspondence $a \leftrightarrow (a_B, a_K)$ is clearly 1-1 and if we assume the connection to be complete, we see that we have built an homeomorphism

$$P(A_E) \rightarrow P(A_B) \times_B P(\mathcal{K}),$$

where $P(A_B) \times_B P(\mathcal{K})$ denotes couples $(a_B, a_K) \in P(A_B) \times P(\mathcal{K})$ such that a_K is a $\mathcal{K}$ -path lying over $\gamma_B(0)$. Recall that $P(A_E)$ enjoys a natural structure of a Banach space (see [6] for more details)

Now if we look at how concatenation in $P(A_E)$ behaves through this homeomorphism, we see that the concatenation $v \cdot u$ of two A_E -paths $u, v \in P(A_E)$ is sent to $(v_B \cdot u_B, \Phi_{u_B}^{-1}(v_K) \cdot u_K)$ where we have set:

$$(18) \quad \Phi_{u_B}^{-1}(v_K)(t) := \Phi_{0,1}^{u_B} \circ v_K(t).$$

Thus we have proved the following:

Proposition 4.1. (*Splitting A_E -paths*). *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a Lie algebroid extension, and fix an Ehresmann connection which is assumed to be complete. Then there is a homeomorphism:*

$$\begin{array}{ccc} P(A_E) & \rightarrow & P(A_B) \times_B P(\mathcal{K}) \\ a & \mapsto & (a_B, a_K). \end{array}$$

under which the concatenation in $P(A_E)$ writes:

$$(19) \quad (a_B^2, a_K^2) \cdot (a_B^1, a_K^1) = (a_B^2 \cdot a_B^1, \Phi_{a_B^1}^{-1}(a_K^2) \cdot a_K^1).$$

Remark 4.2. Let us explain the picture we have in mind: given an algebroid $A \rightarrow M$, we can think of $P(A)$ as a “groupoid” over M : source and target are defined the obvious way and concatenation plays the role of multiplication. One also has the notions of “units” and of “inverses”: $1_m(t) := 0_m$, and $a^{-1}(t) := -a(1-t)$ (note however that $a \cdot a^{-1} \neq 1_{\gamma(0)}$). Morphisms of “groupoids” should only commute with source, target and multiplication, and sent units on units, the most obvious way to obtain such a morphism being induced by a algebroid morphism (rather than introducing useless vocabulary, we just stick to “groupoids” without further precisions).

Generalizing the notion of action introduced in Appendix B to “groupoids”, we can use the holonomy $\Phi : P(A_B) \mapsto \text{Gauge}(P(\mathcal{K}))$ to get an action of $P(A_B) \rightrightarrows B$ on the fibered “groupoid” $P(\mathcal{K}) \rightarrow B$, and then build the corresponding action “groupoid” $P(A_B) \times_B P(\mathcal{K}) \rightrightarrows E$. Then we see that the formula for composition is exactly the one given in the Proposition 4.1. Thus we can think of $P(A_E)$ as a groupoid associated to an action on a fibered groupoid.

Because of the presence of curvature, there is no action of the groupoid $\mathcal{G}(A_B)$ neither on $P(\mathcal{K})$, nor on $\mathcal{G}(\mathcal{K})$. However, it is still possible to describe

A_E -homotopies through this homeomorphism in a reasonable way, as stated in next proposition.

Let us before precise a natural procedure to extend a A_E -path a into a time-dependant section of A_E we will always implicitly use: first extend arbitrarily $a_B(t)$ into $\alpha_B(t) \in \Gamma(A_B)$ and a_V into a time-dependant section α_V of $\mathcal{K} \subset A_E$. Clearly, $\alpha := h(\alpha_B(t)) + \alpha_V$ extends a . Moreover, $\alpha_K(t) := (\Phi_{0,t}^{a_B})_*(\alpha_V(t))$ extends the $\mathcal{K}$ -path a_K defined by (17).

Proposition 4.3. *Let a^ϵ be a smooth family of A_E -paths and $(a_B^\epsilon, a_K^\epsilon)$ the corresponding family in $P(A_B) \ltimes P(\mathcal{K})$ given by last proposition.*

Then a^ϵ defines a A_E -homotopy if and only a_B^ϵ is a A_B -homotopy such that the unique solution $\mu_K \in \Gamma(\mathcal{K})$ of the evolution equation:

$$(20) \quad \frac{d\alpha_K}{d\epsilon} - \frac{d\mu_K}{dt} = [\alpha_K, \mu_K]_{\mathcal{K}},$$

with initial condition $\mu_K^0(\epsilon) = 0$ satisfies:

$$(21) \quad \mu_K^1(\epsilon)_{\gamma_K^\epsilon(1)} = \int_0^1 (\Phi_{0,s}^{a_B^\epsilon})_* \left(\omega(a_B, b_B)_{s,\epsilon} \right) ds, \quad \forall \epsilon \in I,$$

Here, $a_B dt + b_B d\epsilon : TI^2 \rightarrow A_B$ is the Lie algebroid morphism induced by the A_B -homotopy a_B^ϵ , and α_K is extending a_K as explained above.

Proof. We first extend the smooth family of A_E -paths a^ϵ into a smooth family of time-dependant sections α^ϵ as specified above: $\alpha = h(\alpha_B) + \alpha_V$. It is easily checked that the solution β of the evolution equation (33) is π -projectable provided α and the initial condition are. Thus, we can write the solution as $\beta = h(\beta_B) + \beta_V$ and use the formula for brackets (9) (10) (11). We see that the evolution equation (33) in A_E is equivalent to the following two evolution equations

$$(22) \quad \frac{d\alpha_B}{d\epsilon} - \frac{d\beta_B}{dt} = [\alpha_B, \beta_B]_{A_B},$$

$$(23) \quad \mathcal{D}_{\beta_B}(\alpha_V) + \frac{d\alpha_V}{d\epsilon} - \mathcal{D}_{\alpha_B}(\beta_V) + \frac{d\beta_V}{dt} = [\alpha_V, \beta_V]_{\mathcal{K}} + \omega(\alpha_B, \beta_B),$$

with initial conditions $\beta_B^0(\epsilon) = 0$ and $\beta_V^0(\epsilon) = 0$. The homotopy condition (35) is clearly equivalent to $\beta_B^1(\epsilon) = 0, \beta_V^1(\epsilon) = 0$ and in particular, one gets from (22) that a_B^ϵ is an homotopy in A_B .

Let us now look at equation (23): in order to get an evolution equation involving α_K^ϵ , we push-forward (23) by means of $\Phi_{0,t}^{a_B^\epsilon}$:

$$\begin{aligned} & (\Phi_{0,t}^{a_B^\epsilon})_* \left(\mathcal{D}_{\beta_B}(\alpha_V) + \frac{d\alpha_V}{d\epsilon} \right) - (\Phi_{0,t}^{a_B^\epsilon})_* \left(\mathcal{D}_{\alpha_B}(\beta_V) + \frac{d\beta_V}{dt} \right) \\ & = (\Phi_{0,t}^{a_B^\epsilon})_* ([\alpha_V, \beta_V]_{\mathcal{K}} + \omega(\alpha_B, \beta_B)). \end{aligned}$$

Only the first term on the left-hand side of this equality is problematic, it will be taken care of in the next lemma. This lemma shows that (23) holds

if and only if the following holds:

$$\begin{aligned} & \frac{d}{d\epsilon}(\Phi_{0,t}^{a_B^\epsilon})_*(\alpha_V) - \left[\int_0^t (\Phi_{0,s}^{a_B^\epsilon})_*(\omega(\alpha_B, \beta_B)_{s,\epsilon}) ds, (\Phi_{0,t}^{a_B^\epsilon})_*(\alpha_V) \right]_{\mathcal{K}} \\ & - \frac{d}{dt}(\Phi_{0,t}^{a_B^\epsilon})_*(\beta_V) = [(\Phi_{0,t}^{a_B^\epsilon})_*(\alpha_V), (\Phi_{0,t}^{a_B^\epsilon})_*(\beta_V)]_{\mathcal{K}} + (\Phi_{0,t}^{a_B^\epsilon})_*(\omega(\alpha_B, \beta_B)). \end{aligned}$$

In other words, if we set

$$\begin{cases} \beta_K^t(\epsilon) &:= (\Phi_{0,t}^{a_B^\epsilon})_*(\beta_V^t(\epsilon)), \\ F_{\epsilon,t} &:= \int_0^t (\Phi_{0,s}^{a_B^\epsilon})_*(\omega(\alpha_B, \beta_B)_{s,\epsilon}) ds, \end{cases}$$

we see that (23) holds if and only if $\beta_K + F$ is solution of equation (20):

$$\frac{d}{dt}(\beta_K + F) - \frac{d\alpha_K}{d\epsilon} + [\alpha_K, \beta_K + F]_{\mathcal{K}} = 0.$$

Now observe that the homotopy condition $\beta_V^1(\epsilon) = 0$ holds if and only if $\beta_K^1(\epsilon) = 0$ (here, we voluntarily omit base points). By uniqueness of the solution $\mu_K = \beta_K + F$ of (20) we see that $\beta_V^1(\epsilon) = 0$ if and only if (21) holds. $\square$

Lemma 4.4. *With the same notations as above, one has:*

$$(\Phi_{0,t}^{a_B^\epsilon})_*(\mathcal{D}_{\beta_B}(\alpha_V) + \frac{d\alpha_V}{d\epsilon}) = \frac{d}{d\epsilon}(\Phi_{0,t}^{a_B^\epsilon})_*(\alpha_V) - [F, (\Phi_{0,t}^{a_B^\epsilon})_*(\alpha_V)]_{\mathcal{K}}.$$

Proof. Consider the solution σ of the following evolution equation:

$$(24) \quad \frac{d\sigma}{dt} - \frac{dh(\alpha_B)}{d\epsilon} + [h(\alpha_B), \sigma]_{A_E} = 0,$$

with initial condition $\sigma^0(\epsilon) = 0$. We can use the same principle as in the proof of Proposition A.1 to show that the flow ψ^{σ^t} of the time-dependant derivation $ad_{\sigma^t}^{A_E}$ satisfies:

$$(25) \quad \psi_{0,\epsilon}^{\sigma^t} = \Phi_{t,0}^{a_B^0} \circ \Phi_{0,t}^{a_B^\epsilon}.$$

Let us now relate σ and $h(\beta_B)$: first applying $h(-)$ to (22), one gets:

$$\frac{dh(\beta_B)}{dt} - \frac{dh(\alpha_B)}{d\epsilon} + h([\alpha_B, \beta_B]_{A_B}) = 0.$$

In this equality, one can subtract equation (24) to get the following:

$$(26) \quad \frac{d(\sigma - h(\beta_B))}{dt} + [h(\alpha_B), \sigma - h(\beta_B)]_{A_E} = \omega(\alpha_B, \beta_B).$$

By applying $(\Phi_{0,t}^{a_B^\epsilon})_*$ to this last equation, we obtain the desired relation between σ and $h(\beta_B)$:

$$(27) \quad \frac{d}{dt}(\Phi_{0,t}^{a_B^\epsilon})_*(\sigma - h(\beta_B)) = (\Phi_{0,t}^{a_B^\epsilon})_*(\omega(\alpha_B^{\epsilon,t}, \beta_B^{\epsilon,t})).$$

We now have enough material to prove the equality stated by the lemma: decompose the left-hand term as follows:

$$(\Phi_{0,t}^{a_B^\epsilon})_*(\mathcal{D}_{\beta_B}(\alpha_V) + \frac{d\alpha_V}{d\epsilon}) = (\Phi_{0,t}^{a_B^\epsilon})_*\left([\sigma, \alpha_V] + \frac{d\alpha_V}{d\epsilon}\right) + (\Phi_{0,t}^{a_B^\epsilon})_*([h(\beta_B) - \sigma, \alpha_V]).$$

In this expression, we first check using (25) that the first term gives $\frac{d}{d\epsilon}(\psi_{0,t}^{\gamma_B^\epsilon})_*(\alpha_V)$:

$$\begin{aligned} (\Phi_{0,t}^{a_B^\epsilon})_*([\sigma, \alpha_V] + \frac{d\alpha_V}{d\epsilon}) &= (\Phi_{0,t}^{a_B^0})_* \circ (\psi_{0,\epsilon}^{\sigma^t})_*([\sigma, \alpha_V] + \frac{d\alpha_V}{d\epsilon}) \\ &= (\Phi_{0,t}^{a_B^0})_*\left(\frac{d}{d\epsilon}(\psi_{0,\epsilon}^{\sigma^t})_*(\alpha_V)\right) \\ &= \frac{d}{d\epsilon}(\Phi_{0,t}^{a_B^0})_* \circ (\psi_{0,\epsilon}^{\sigma^t})_*(\alpha_V) \\ &= \frac{d}{d\epsilon}(\Phi_{0,t}^{a_B^\epsilon})_*(\alpha_V). \end{aligned}$$

It is easily integrated relation (27) to show that the second term gives

$$(\Phi_{0,t}^{a_B^\epsilon})_*([h(\beta_B) - \sigma, \alpha_V]) = [F, (\psi_{0,t}^{\gamma_B^\epsilon})_*(\alpha_V)]_{\mathcal{K}},$$

which completes the proof. $\square$

Now that we have translated the homotopy condition to a condition on the split paths, the theorem below follows easily from Proposition A.1.

Theorem 4.5. *Consider a Lie algebroid extension $K \hookrightarrow A_E \twoheadrightarrow A_B$, endowed with a Ehresmann connection $A_E = \mathcal{K} \oplus H$, which we assume to be complete.*

Then the topological source simply connected groupoid $\mathcal{G}(A_E)$ integrating A_E is naturally identified with the quotient

$$P(A_B) \times_B \mathcal{G}(\mathcal{K}) / \sim$$

where the equivalence relation is given by: $(a_0, g_0) \sim (a_1, g_1)$ if and only if there exists a A_B -homotopy $h_B = \text{adt} + b d\epsilon : TI^2 \mapsto A_B$ between a_0 and a_1 such that:

$$g_1 \cdot g_0^{-1} = \partial(h_B, \mathbf{t}(g_0)).$$

Here, $\partial(h_B, x_0)$ is the element in $\mathcal{G}(\mathcal{K})$ represented by the $\mathcal{K}$ -path:

$$(28) \quad \epsilon \rightarrow \int_0^1 (\Phi_{0,s}^{a_B^\epsilon})_* \left(\omega(a, b)_{s,\epsilon} \right) ds \in \mathcal{K}_{\gamma_K^\epsilon},$$

where $\gamma_K^\epsilon := \phi_{a_B^\epsilon}^{-1} \circ \phi_{a_B^0}(x_0)$.

Example 4.6. In the case of central extensions, Theorem 4.5 reduces to a result due to Crainic in a Lie algebroid approach to prequantization [5], see also [8] in the context of Poisson and Jacobi manifolds. Recall that $E = B$ and $\mathcal{K} := \mathbb{R}^n \times B$ is a bundle of abelian Lie algebras on which A_B acts trivially, while ω is a d_A -closed 2-form on A_B with values in $\mathbb{R}^n$.

Then for any A_B -path a_B covering γ_B , the holonomy $\Phi_{a_B} : \mathcal{K}_{\gamma_B(0)} \rightarrow \mathcal{K}_{\gamma_B(1)}$ is trivial: $\Phi_{a_B}(r, \gamma_B(0)) = (r, \gamma_B(1))$. Moreover $\mathcal{G}(\mathcal{K})$ is identified with the bundle of abelian Lie groups $B \times \mathbb{R}^n$ by averaging $\mathcal{K}$ -paths. This process makes appear a second integral in the formula (28) and the homotopy condition reads as follows: two elements (a_0, g_0) and (a_1, g_1) in $\mathbb{R}^n \times P(A_B)$ are homotopic if and only if there exists a A_B -homotopy $\gamma_B : TI^2 \mapsto A_B$ between γ_0 and γ_1 such that:

$$g_1 - g_0 = \int_{\gamma_B} \omega.$$

The Theorem 4.5 might be seen as a non abelian generalization of this result.

Remark 4.7. There is an obvious way to concatenate two A_B -homotopies $h = adt + bde, \tilde{h} = \tilde{a}dt + \tilde{b}de$ such that $h^{\epsilon=1} = \tilde{h}^{\epsilon=0}$. This amounts to concatenate two algebroid morphisms $TI^2 \rightarrow A_B$ along the edges $\{\epsilon = 1\}$ and $\{\epsilon = 0\}$ of two copies of I^2 .

To do this properly, one needs to reparametrize in the ϵ variable in order to make sure the concatenation is smooth (on the way one sees that it is enough to assume that $a_{\epsilon=1} = \tilde{a}_{|\epsilon=0}$ to be able to concatenate). We leave the details to the reader since the construction is a straightforward adaptation of the concatenation for A -paths as given in [6].

Then it is not hard to see that the operator ∂ defined in the theorem satisfies by construction:

$$\partial(\tilde{h} \cdot h) = \partial\tilde{h} \cdot \partial h.$$

This suggests one should think of A -homotopies as a “groupoid” over $P(A)$. In fact A -homotopies modded out by higher homotopies (so that the concatenation is associative) form a Banach groupoid over $P(A)$, and this groupoid presents a finite dimensional differentiable stack $\mathcal{G}(A)$. As shown in [17], $\mathcal{G}(A)$ is itself again a stacky groupoid with the same base as A and it is in the sense of differentiable stacks the universal integrating object of A .

Remark 4.8. The reader might be surprised that the formula (28) does not involve the values of $\omega(a, b)_{s, \epsilon}$ at the base points $\gamma^\epsilon(s)$ but at $\phi_{s,1}^{a^\epsilon}(\gamma^\epsilon(1))$. Let us explain this fact: for each ϵ , there is an A_E -homotopy between a^ϵ and the concatenation $h(a_B^\epsilon) \cdot a_K^\epsilon$ (this can be proved using Proposition A.1). Here $h(a_B^\epsilon)$ is the only horizontal A_E -path over a_B^ϵ that can be concatenated with a_K^ϵ that is, $h(a_B)$ has base path $s \rightarrow \phi_{s,0}^{a^\epsilon}(\gamma_K^\epsilon(1))$ which precisely coincides with $\phi_{s,1}^{a^\epsilon}(\gamma^\epsilon(1))$ by construction.

Thus we see that the homotopy condition we get in Theorem 4.5 relies on the representatives up to A_E -homotopy $h(a_B^\epsilon) \cdot a_K^\epsilon$ of a^ϵ , rather than on a^ϵ itself.

4.2. The Monodromy Groupoid. We now turn to the study of the sequence of groupoids

$$(29) \quad 1 \rightarrow \mathcal{G}(\mathcal{K}) \xrightarrow{\tilde{i}} \mathcal{G}(A_E) \xrightarrow{\tilde{\pi}} \mathcal{G}(A_B) \rightarrow 1.$$

obtained by integration (see [14], [15]) of a Lie algebroid extension

$$(30) \quad \mathcal{K} \xhookrightarrow{i} A_E \twoheadrightarrow^{\pi} A_B.$$

Recall that $\tilde{i}$ and $\tilde{\pi}$ are defined at the level of paths, as follows:

$$\begin{aligned} \tilde{i}([a_K]_{\mathcal{K}}) &:= [i \circ a_K]_{A_E}, \\ \tilde{\pi}([a]_{A_E}) &:= [\pi \circ a]_{A_B}, \end{aligned}$$

for any $\tilde{a}_K \in P(\mathcal{K})$, and $a \in P(A)$. Our aim in this section is to see why the sequence of groupoids (29) might not be exact in general. In other words, we want to explain the lack of exactness of the *integration functor*. We start by showing that $\tilde{\pi}$ is surjective provided there is a complete Ehresmann connection.

Proposition 4.9. *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a Lie algebroid extension that admits a complete Ehresmann connection. Then the sequence of groupoids (29) is exact at $\mathcal{G}(A_B)$.*

Proof. We have to prove that $\tilde{\pi}$ is surjective. Given any A_B -path $a_B \in P(A_B)$ covering γ_B in B , one extends a_B into a time-dependant section α_B of A_B , and consider its horizontal lift $h(\alpha_B)$. Pick an arbitrary point e_0 in the fiber over $\gamma_B(0)$, then clearly $a(t) := h(\alpha_B)_{\phi_{t,0}^{a_B}(e_0)}$ defines a A_B -path that projects onto a_B . Completeness of the Ehresmann connection insures us that a is well defined for all $t \in I$. $\square$

Remark 4.10. We stress the fact that completeness is needed in order to ensure surjectivity, otherwise one might only get a quasi-surjective morphism in the sense that for any $x \in \mathcal{G}(A_B)$, there exists $g_1, \dots, g_n \in \mathcal{G}(A_E)$ such that the composition $\tilde{\pi}(g_1) \cdots \tilde{\pi}(g_n) = x$. In general though, the g_i 's might not be chosen to be composable, as the following examples from [20] show.

Example 4.11. (*Stairway To Heaven*). Consider $E = \{(x, y) \in \mathbb{R}^2, y \in \mathbb{Z}, x \in]y-1, y+1[\}$, it is a smooth submanifold of $\mathbb{R}^2$, and the projection on the first factor is a submersion onto $B := \mathbb{R}$. Thus one gets a well defined algebroid morphism $TE \rightarrow TB$, that comes with a unique Ehresmann connection. It is easily seen that the induced groupoid morphism is only quasi-surjective.

Example 4.12. (*Highway To Hell*). Consider now on $E = \mathbb{R}^2$ the integrable regular foliation $\mathcal{F} \subset TE$ spanned by $\partial x - \exp(y)\partial y$. Denote p the projection on the first factor. The restriction of dp to the Lie algebroid $A_E := \mathcal{F}$ defines a surjective Lie algebroid morphism onto $A_B := TB \rightarrow B := \mathbb{R}$. Again, we have a (unique) Ehresmann connection which is not complete: $h(\partial x) = \partial x - \exp(y)\partial y$. Lie algebroid morphism $dp|_{\mathcal{F}}$ only integrates into a quasi-surjective Lie groupoid morphism.

Proposition 4.13. *Let $\mathcal{K} \hookrightarrow A_E \rightrightarrows A_B$ be a Lie algebroid extension that admits a complete Ehresmann connection. Then the sequence of groupoids (29) is exact at $\mathcal{G}(A_E)$.*

Proof. By definition, elements in $\ker \tilde{\pi}$ are represented by A_E -paths whose projection on A_B is homotopic to a trivial path. In particular, an element of the form $i \circ a_K$ projects onto a trivial A_B -path since $\pi \circ i \circ a_K = 0$. Thus we have $\text{Im } \tilde{\iota} \subset \ker \tilde{\pi}$ in a (very) trivial way.

Conversely, consider any element of $\ker \tilde{\pi}$ represented by some A_E -path a , and extend a into a time-dependant section α as prescribed in last section. With the same notations, we get an identification $a = (a_K, a_B)$ where, by assumption a_B is homotopic to a trivial path by means of some homotopy $h_B = a_B dt + b_B d\epsilon : TI^2 \rightarrow A_B$.

One extends b_B into a family of time-dependant sections β_B of A_B such that $\beta_B|_{t=0,1} = 0$. Thus, if we let α^ϵ be the solution of the evolution equation $\frac{d\alpha}{d\epsilon} - \frac{dh(\beta_B)}{dt} = [\alpha, h(\beta_B)]$ with initial condition $\alpha^{\epsilon=0} := \alpha$, we get a A_E -homotopy $(a_K^\epsilon, a_B^\epsilon)$. By construction $(a_K^1, a_B^1) = (a_K^1, 0)$ which clearly represents an element in $\text{Im } \tilde{\iota}$. Thus we have shown that any element of $\ker \tilde{\pi}$ is represented by a A_E -path of the form $(a_K, 0_x)$, thus it is an element of $\text{Im } \tilde{\iota}$. $\square$

We now see that, when there is a complete connection, the lack of exactness in the sequence 29 can only occur at $\mathcal{G}(\mathcal{K})$. This means that $\mathcal{G}(\mathcal{K})$

might not inject into $\mathcal{G}(A_E)$. In order to measure this, we introduce the following.

Definition 4.14. We will call **monodromy groupoid** the kernel of $\tilde{\iota}$, and denote it $\mathcal{M}$.

We obtain by construction an exact sequence of groupoids:

$$(31) \quad 1 \rightarrow \mathcal{M} \rightarrow \mathcal{G}(\mathcal{K}) \rightarrow \ker \tilde{\pi} \rightarrow 1.$$

The next theorem explains how $\mathcal{M}$ is involved in the homotopy theories of A_E , A_B and $\mathcal{K}$. In fact, it is the image of a connecting morphism $\partial_2 : \pi_2(A_B) \times_B E \rightarrow \mathcal{G}(\mathcal{K})$, where $\pi_2(A_B)$ denote the homotopy classes of spheres in A_B :

Theorem 4.15. *Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a Lie algebroid extension that admits a complete Ehresmann connection. Then there exists a homomorphism*

$$\partial_2 : \pi_2(A_B) \times_B E \rightarrow \mathcal{G}(\mathcal{K}),$$

that makes the following sequence exact:

$$\cdots \rightarrow \pi_2(A_B) \times E \xrightarrow{\partial_2} \mathcal{G}(\mathcal{K}) \xrightarrow{\tilde{\iota}} \mathcal{G}(A_E) \twoheadrightarrow \mathcal{G}(A_B)$$

Proof. The boundary map ∂_2 can be constructed directly as follows: consider an element x_0 in E , b_0 its projection onto B , and s_B a A_B -sphere based at b_0 i.e. an algebroid morphism $s_B = a_B dt + b_B d\epsilon : I \times I \rightarrow B$ satisfying $a_B|_{\epsilon=0,1} = 0$, $b_B|_{t=0,1} = 0$ (see the appendix C).

Choose any family β_B^t of time-dependant sections of A_B extending b_B , chosen such that $\beta_B^{t=0} = \beta_B^{t=1} = 0$. Thus, the unique solution α_B of the evolution equation $[h(\beta_B), \alpha]_{A_E} = \frac{d}{dt}h(\beta_B) - \frac{d}{d\epsilon}\alpha$ with initial condition $\alpha^{\epsilon=0} = 0$ induces at time $\epsilon = 1$ a $\mathcal{K}$ -path starting at x_0 , which is homotopic (as a A_E -path) to the trivial path 0_{x_0} by construction. It is easily seen using the construction in the proof of Proposition 4.3 that a^1 is a $\mathcal{K}$ -path.

One defines $\partial_2(h_B, x_0)$ to be the $\mathcal{K}$ -homotopy class of a^1 . Clearly, $\partial_2(h_B, x_0)$ has source and target x_0 , so it is an element in the isotropy $\mathcal{G}(\mathcal{K})_{x_0}$.

We now show that $\partial(h_B, x_0)$ only depends on the homotopy class of s_B . Consider an homotopy of A_B -spheres $s_B^u : TI^2 \rightarrow A_B, u \in I$ between two spheres $s_B^{u=0}$ and $s_B^{u=1}$; we just do the same construction as above with u as a parameter: we extend b_B^u into a smooth family β_B^u of sections of A_B vanishing whenever $t = 0, 1$ and we let $\alpha^{\epsilon,u}(t)$ be the unique solution of

$$\frac{d}{dt}\alpha - \frac{d}{d\epsilon}h(\beta_B) = [h(\beta_B), \alpha]_{A_E},$$

with initial condition $\alpha|_{\epsilon=0} = 0$. We let $a_K^{u,\epsilon=1}$ the corresponding $\mathcal{K}$ -path.

We now want to define a $\mathcal{K}$ -homotopy between $a_K^{u=0,\epsilon=1}$ and $a_K^{u=1,\epsilon=1}$. For this, we consider the unique solution of the evolution equation:

$$\frac{d\theta}{d\epsilon} - \frac{dh(\beta_B)}{du} = [\theta, h(\beta_B)]_{A_E},$$

with initial condition $\theta|_{\epsilon=0} = 0$. We know from the Lemma C.6 that θ also satisfies:

$$\frac{d\theta}{dt} - \frac{d\alpha}{du} = [\theta, \alpha]_{A_E}.$$

and that $\theta|_{t=0,1} = 0$ so it gives the desired homotopy.

To complete the proof, one only needs to apply theorem 4.5 to see that every element in $\mathcal{M}$ is obtained this way. $\square$

Remark 4.16. The operator defined above clearly coincides with the one defined in Theorem 4.5 applied to A -spheres. In fact, the arguments in the proof above show that $\partial(h) = \partial(\hat{h})$ provided h and $\hat{h}$ are equivalent homotopies between two fixed A_B -paths a_B^0 and a_B^1 . By equivalent homotopies, we mean by definition that there exists a Lie algebroid morphism $h =adt + bde + cdu : TI^3 \rightarrow A$ with $a|_{u=0}dt + b|_{u=0}de = h$, $a|_{u=1}dt + b|_{u=1}de = \hat{h}$ and satisfying $c = 0$ whenever e or t is in $\{0, 1\}$.

4.3. Integrability. The analogy between the monodromy groupoid described above and the monodromy groups of a Lie algebroid as constructed in [6] is clear. In fact it is easily seen that they do coincide in the case of Atiyah exact sequences. Also, when E is a tubular neighborhood of an orbit B of A_E (see Example 2.20), the restriction of the monodromy groupoid $\mathcal{M}$ to B coincides with the monodromy (bundle of) groups along B by construction.

Just like monodromy groups control the integrability of a Lie algebroid ([6]), in the case of clean extensions (see the Definition 2.9) the monodromy groupoid controls the integrability of A_E .

Definition 4.17. we will say that $\mathcal{M}$ is discrete near identities if, for any sequence $(g_n)_n \subset \mathcal{M}$ converging to an identity $\mathbf{1}_x$ in $\mathcal{G}(\mathcal{K})$, we have $g_n = \mathbf{1}_{x_n}$ for n big enough.

Theorem 4.18. Let $\mathcal{K} \hookrightarrow A_E \twoheadrightarrow A_B$ be a clean Lie algebroid extension. Assume that it admits a complete Ehresmann connection and that both $\mathcal{K}$ and A_B are integrable Lie algebroids.

Then A_E is integrable if $\mathcal{M}$ is discrete near identities in $\mathcal{G}(\mathcal{K})$.

Proof. For a generic Lie algebroid $A \rightarrow M$, we denote:

$$\mathcal{N}_x^A := \{[g] \in \mathcal{G}(\mathfrak{g}_x^A), g \text{ is } A\text{-homotopic to a trivial path}\},$$

where $g \in P(\mathfrak{g}_x^A)$ and $x \in M$. According to [6] we have an exact sequence:

$$\mathcal{N}_x^A \hookrightarrow \mathcal{G}(\mathfrak{g}_x^A) \rightarrow \mathcal{G}(A)_x^0.$$

Moreover, any element in $\mathcal{N}_x(A)$ can be represented by an element in $Z(\mathfrak{g}_x^A)$, seen as a constant path (here, $Z(\ker \sharp_{Ax})$ denotes the center of the isotropy algebra $\mathfrak{g}_x^A$ at x). With these settings, integrability is equivalent to the following condition:

For any sequence $([v_n])$ in $\mathcal{N}_{x_n}^A$ represented by constant paths $v_n \in Z(\ker \sharp_{Ax_n})$ that converges to a trivial path 0_x in $P(A)$, one necessarily has $v_n = 0_{x_n}$ for n big enough.

So let $([v_n]) \subset \mathcal{N}_{x_n}^{A_E}$ be a sequence where $v_n \in Z(\mathfrak{g}_{x_n}^E)$ converges to a trivial path: $v_n \rightarrow 0_x$ for some $x \in E$. For each n , one can draw the following

commutative diagram:

$$\begin{array}{ccccc}
\mathcal{N}_{x_n}^{\mathcal{K}} & \hookrightarrow & \mathcal{G}(\mathfrak{g}_{x_n}^{\mathcal{K}}) & \twoheadrightarrow & \mathcal{G}(\mathcal{K})_{x_n}^0 \\
& & \downarrow \tilde{\iota} & & \downarrow \\
\mathcal{N}_{x_n}^{A_E} & \hookrightarrow & \mathcal{G}(\mathfrak{g}_{x_n}^E) & \twoheadrightarrow & \mathcal{G}(A_E)_{x_n}^0 \\
& & \downarrow \tilde{\pi} & & \downarrow \\
\mathcal{N}_{y_n}^{A_B} & \hookrightarrow & \mathcal{G}(\mathfrak{g}_{y_n}^B) & \twoheadrightarrow & \mathcal{G}(A_B)_{y_n}^0,
\end{array}$$

where $y_n := p(x_n) \in B$. As explained above all lines are exact. The middle column is not exact but we still have $\text{Im } \tilde{\iota} = \ker \tilde{\pi}$.

Now consider the sequence of A_B -paths $\pi(v_n)$ in $\mathfrak{g}_{y_n}^B$. The restriction of π to $Z(\mathfrak{g}_{x_n}^E)$ is easily seen to induce an exact sequence

$$(32) \quad Z(\mathfrak{g}_{x_n}^{\mathcal{K}}) \hookrightarrow Z(\mathfrak{g}_{x_n}^E) \rightarrow Z(\mathfrak{g}_{y_n}^B).$$

Thus $\tilde{\pi}([v_n]) = [\pi(v_n)]$ is a sequence in $\mathcal{N}_{x_n}^{A_B}$ with $\pi(v_n) \in Z(\ker \sharp_{A_B y_n})$ converging to 0_{y_n} . Since A_B is integrable, this implies that $\pi(v_n) = 0_{y_n}$ for n big enough. So we see in (32) that $v_n \in Z(\mathfrak{g}_{x_n}^{\mathcal{K}})$ and converges to 0_{x_n} . By construction, we have $[v_n]_{\mathcal{K}} \in \mathcal{M}$ and converges to the identity $\mathbf{1}_x$ in $\mathcal{G}(\mathcal{K})$. The discreteness assumption on $\mathcal{M}$ implies that $[v_n]_{\mathcal{K}} = \mathbf{1}_{x_n}$ for n big enough. Thus $[v_n] \in \mathcal{N}_{x_n}^{\mathcal{K}}$ with $v_n \in Z(\mathfrak{g}_{x_n}^{\mathcal{K}})$ converging to 0_x . We can conclude that $v_n = 0_{x_n}$ for n big enough because $\mathcal{K}$ is integrable. $\square$

APPENDIX A. THE WEINSTEIN GROUPOID AND A -HOMOTOPIES

Let $A \rightarrow M$ be a Lie algebroid. We recall the construction of the Weinstein groupoid $\mathcal{G}(A)$ (see [6] or [4] for an alternative approach in the Poisson case). This is a topological groupoid, with source 1-connected fibers, which morally integrates A . Indeed, A is integrable if and only if $\mathcal{G}(A)$ is smooth and in this case $A(\mathcal{G}(A))$ is canonically isomorphic to A .

We will denote by $P(A)$ the space of A -paths (up to reparametrization). By setting $s(a) = p_A \circ a(0)$ and $t(a) = p_A \circ a(1)$, we shall think of $P(A) \rightrightarrows M$ as an infinite dimensional groupoid with multiplication given by concatenation (though units are not well defined). On the groupoid $P(A)$ there is an equivalence relation $\sim$, called A -homotopy, which preserves products, and one sets:

$$\mathcal{G}(A) := P(A) / \sim.$$

The homotopy class of an A -path a will be denoted $[a]_A$ or $[a]$ when no confusion seems possible. Let us recall how A -homotopies are defined since this will be essential later. Suppose we are given $\alpha^\epsilon(t)$, a time dependent family of sections of A depending on a parameter $\epsilon \in I := [0, 1]$, and $\beta^0(\epsilon)$ a time dependent section of A . Then there exists a unique solution $\beta = \beta^t(\epsilon)$ of the following *evolution equation*:

$$(33) \quad \frac{d\alpha}{d\epsilon} - \frac{d\beta}{dt} = [\alpha, \beta],$$

with initial condition $\beta^0(\epsilon)$. In fact, it is easily checked that the following integral formula provides a solution:

$$(34) \quad \beta^t(\epsilon) := \int_0^t (\psi_{t,s}^{\alpha^\epsilon})_* \left(\frac{d}{d\epsilon} \alpha^\epsilon(s) \right) ds + (\psi_{t,0}^{\alpha^\epsilon})_*(\beta^0(\epsilon)).$$

Here ψ^{α^ϵ} denotes the flow of the time-dependant linear vector field on A , associated to the derivation $[\alpha^\epsilon, -]$ of sections of A (see the appendix in [6] for more details). We emphasize the use of the indices and parameters in the notation: we think of α as an ϵ -family of t -time dependent sections of A , while we think of β as a t -family of ϵ -time dependent sections of A (see why below).

The notion of A -homotopy is defined as follows. A family $a^\epsilon : I \mapsto A$, $\epsilon \in I$ of A -paths, over $\gamma^\epsilon : I \mapsto B$ is called a *homotopy* if $\gamma^\epsilon(0)$ is independent of ϵ , and if the unique solution β of equation (33) with initial condition $\beta^0(\epsilon) = 0$ satisfies:

$$(35) \quad \beta^1(\epsilon)_{\gamma^\epsilon(1)} = 0, \forall \epsilon \in I.$$

Here, α^ϵ denotes any family of time-dependant sections of A extending a , that is, such that $\alpha_{\gamma^\epsilon(t)}^\epsilon(t) = a^\epsilon(t)$. One checks that this definition is independent of the choice of α (see [6]). We will refer to (35) as the *homotopy condition*. Note that, if some α^0 is fixed, and we are given β with $\beta^0 = \beta^1 = 0$, then equation (33) can also be considered as an evolution equation for α , which turns out to induce an homotopy.

The reason why we consider the evolution equation (33) with non-vanishing initial condition is that this also leads to A -homotopies, as we now explain. Set $X = \sharp\alpha$ and $Y = \sharp\beta$, so that X^ϵ and Y^t are families of time dependent vector fields on M (which obviously satisfy an evolution equation in the algebroid TM). As the proof of next proposition shows, this forces their time-dependant flows $\phi_{t,0}^{X^\epsilon}, \phi_{\epsilon,0}^{Y^t}$ to be related as follows:

$$(36) \quad \phi_{t,0}^{X^\epsilon} \circ \phi_{\epsilon,0}^{Y^0} = \phi_{\epsilon,0}^{Y^t} \circ \phi_{t,0}^{X^0}.$$

In particular, if we denote by $\gamma^\epsilon(t)$ any of these two equivalent expressions applied to some $m_0 \in M$, we obtain two families of A -paths a^ϵ and b^t , defined by $a^\epsilon(t) := \alpha^\epsilon(t)_{\gamma^\epsilon(t)}$ and $b^t : \epsilon \mapsto \beta^t(\epsilon)_{\gamma^\epsilon(t)}$. We have:

Proposition A.1. *For any couple α and β satisfying the evolution equation (33), the concatenations $a^1.b^0$ and $b^1.a^0$ defined above are homotopic A -paths.*

Proof. Assume that A is integrable and denote by $\mathcal{G}(A)$ the source 1-connected Lie groupoid integrating A . Recalling that the Lie algebra of sections of A can be identified with the Lie algebra of right invariant vector fields on $\mathcal{G}$, we denote by $\vec{\alpha}^\epsilon$ and $\vec{\beta}_R^t$ the (time dependent) right invariant vector fields on $\mathcal{G}(A)$ that correspond to α^ϵ and β^t , respectively. Then the evolution equation (33) exactly means that $\vec{\alpha}^\epsilon + \partial t$ and $\vec{\beta}^t + \partial \epsilon$ commute, when seen as vector fields on $\mathcal{G}(A) \times I \times I$. This means that their flows:

$$\begin{aligned} \phi_u^{\alpha+\partial t}(g, t, \epsilon) &= (\phi_{t+u,t}^{\vec{\alpha}^\epsilon}(g), t+u, \epsilon) \\ \phi_v^{\beta+\partial \epsilon}(g, t, \epsilon) &= (\phi_{\epsilon+v,\epsilon}^{\vec{\beta}^t}(g), t, \epsilon+v), \end{aligned}$$

commute, so we find:

$$\phi_{t+u,t}^{\vec{\alpha}^{\epsilon+v}} \circ \phi_{\epsilon+v,\epsilon}^{\vec{\beta}^t}(g) = \phi_{\epsilon+v,\epsilon}^{\vec{\beta}^{t+u}} \circ \phi_{t+u,t}^{\vec{\alpha}^\epsilon}(g).$$

In particular, taking $t = \epsilon = 0$ and then switching the roles of u, v with the ones of t, ϵ , one gets:

$$\phi_{t,0}^{\vec{\alpha}^\epsilon} \circ \phi_{\epsilon,0}^{\vec{\beta}^0}(1_{m_0}) = \phi_{\epsilon,0}^{\vec{\beta}^t} \circ \phi_{t,0}^{\vec{\alpha}^0}(1_{m_0}).$$

Now, observe that the path in $\mathcal{G}$ corresponding to the concatenation $a^1.b^0$ is the concatenation of the following paths

$$\epsilon \mapsto \phi_{\epsilon,0}^{\vec{\beta}^0}(1_{m_0}), \quad t \mapsto \phi_{t,0}^{\vec{\alpha}^1} \circ \phi_{1,0}^{\vec{\beta}^0}(1_{m_0}),$$

while the one corresponding to $b^1.a^0$ is the concatenation of the paths

$$t \mapsto \phi_{t,0}^{\vec{\alpha}^0}(1_{m_0}), \quad \epsilon \mapsto \phi_{\epsilon,0}^{\vec{\beta}^1} \circ \phi_{1,0}^{\vec{\alpha}^0}(1_{m_0}).$$

These are clearly homotopic (in the $\mathbf{s}$ -fibers) since they define the boundary of the square:

$$(t, \epsilon) \mapsto \phi_{t,0}^{\vec{\alpha}^\epsilon} \circ \phi_{\epsilon,0}^{\vec{\beta}^0}(1_{m_0}) = \phi_{\epsilon,0}^{\vec{\beta}^t} \circ \phi_{t,0}^{\vec{\alpha}^0}(1_{m_0}).$$

We conclude that $a^1.b^0$ and $b^1.a^0$ are homotopic as A -paths.

In the case A is not integrable, the above proposition still holds. The argument goes as follows: $adt + bde : TI^2 \rightarrow A$ is a Lie algebroid morphism (as can be checked in local coordinates). Let $h : I^2 \rightarrow I^2$ be any homotopy between $u^1 \cdot v^0$ and $v^1 \cdot u^0$, where $u^1 \cdot v^0$ is the concatenation of $v : \epsilon \mapsto (0, \epsilon)$ with $u^1 : t \rightarrow (t, 1)$ and $v^1 \cdot u^0$ the concatenation of $u^0 : t \rightarrow (t, 0)$ with $v^1 : \epsilon \rightarrow (1, \epsilon)$. Then the composition $(adt + bde) \circ dh : TI^2 \rightarrow A$ is a Lie algebroid morphism that defines a homotopy between $a^1 \cdot b^0$ and $b^1 \cdot a^0$. $\square$

Remark A.2. We can also interpret the proposition in terms of A -homotopies (when $b^0 = 0$): it says that b^1 is a representative (up to A -homotopy) of $a^1.(a^0)^{-1}$. Hence, b^1 is trivial if and only if a^0 is homotopic to a^1 .

APPENDIX B. FIBERED LIE GROUPOIDS

We briefly explain basic notions about groupoids acting on groupoids, and how to form semi-direct products.

Definition B.1. Given a submersion $p : E \rightarrow B$, a **fibred Lie groupoid** is a Lie groupoid $\mathcal{G}_V \rightrightarrows E$ over E such that $p \circ \mathbf{s} = p \circ \mathbf{t}$.

Clearly, the orbits of a fibred Lie groupoid lie in the fibers of p , thus the restrictions $\mathcal{G}_V|_{E_b} \rightrightarrows E_b$ are Lie groupoids as well for any $b \in B$.

We will denote $p : \mathcal{G}_V \rightarrow B$ rather than $p \circ \mathbf{s}$ or $p \circ \mathbf{t}$ (there is no real confusion to be done here). It is a Lie groupoid morphism onto the groupoid B (whose only arrows are unities) so one can actually think of $\mathcal{G}_V$ as an extension of B with trivial kernel.

Definition B.2. We will call the **gauge groupoid**, denoted $\text{Gauge}(\mathcal{G}_V) \rightrightarrows B$, the set of all groupoid isomorphisms $\Phi_{b_2, b_1} : \mathcal{G}_V|_{E_{b_1}} \rightarrow \mathcal{G}_V|_{E_{b_2}}$, with source and target $\mathbf{s}(\Phi_{b_2, b_1}) = b_1, \mathbf{t}(\Phi_{b_2, b_1}) = b_2$ and obvious identities and composition.

Of course the gauge groupoid is not Lie in general, however it is a nice intermediary in order to define actions of a Lie groupoid onto an fibered one.

Definition B.3. A smooth **action** of a Lie groupoid $\mathcal{G}_B \rightrightarrows B$ on a fibered Lie groupoid $p : \mathcal{G}_V \rightarrow B$ is a groupoid morphism $\Phi : \mathcal{G}_B \rightarrow \text{Gauge}(\mathcal{G}_V)$ covering the identity and such that

$$\begin{aligned} \mathcal{G}_B \times_B \mathcal{G}_V &\rightarrow \mathcal{G}_V \\ (a_B, a_V) &\rightarrow \Phi_{a_B}(a_V) \end{aligned}$$

is a smooth map. Here, of course $\mathcal{G}_B \times_B \mathcal{G}_V := \{(a_V, a_B) \in \mathcal{G}_B \times \mathcal{G}_V, \mathbf{s}(a_B) = p(a_V)\}$ with the fibered topology.

Given smooth action of a Lie groupoid $\mathcal{G}_B \rightrightarrows B$ on a fibered Lie groupoid $p : \mathcal{G}_V \rightarrow B$ there is a natural structure of Lie groupoid on $\mathcal{G}_B \times_B \mathcal{G}_V \rightarrow E$. Source and target are given by $\mathbf{s}(a_B, a_V) = \mathbf{s}(a_V)$, $\mathbf{t}(a_B, a_V) = \mathbf{t}(\Phi_{a_B}(a_V))$ and composition by:

$$(a_B^2, a_V^2) \cdot (a_B^1, a_V^1) = (a_B^2 \cdot a_B^1, \Phi_{a_B^1}^{-1}(a_V^2) \cdot a_V^1).$$

We will leave the details to the reader. Just note that this is a bit different from the usual notion of action of a groupoid on a submersion $\mathcal{G}_V \rightarrow E$ since the resulting groupoid is over E rather than $\mathcal{G}_V$.

APPENDIX C. SPHERES IN A LIE ALGEBROID

We want to extend the notion of homotopy groups to algebroids. In the case of an integrable algebroid $A \rightarrow M$ these should coincide with the second homotopy groups of the $\mathbf{s}$ -fibers, seen as a completely intransitive groupoid over M , however one needs a construction which is independent of integrability.

Definition C.1. For a Lie algebroid A , we define a **A -sphere** to be an algebroid morphism $s = adt + bde : TI^2 \rightarrow A$ such that $a|_{t=0,1}$ and $b|_{t=0,1}$ vanish (i.e. $a : I^2 \rightarrow A$ (resp. $b : I^2 \rightarrow A$) vanishes whenever $t \in \{0, 1\}$ (resp. $e \in \{0, 1\}$). The space of all such spheres will be denoted $S^2(A)$.

Clearly, a A -sphere $s : TI^2 \rightarrow A$ covers a topological sphere in the base manifold, that is, the base map of s is a map $\gamma : I^2 \rightarrow M$ whose restriction to the boundary of I^2 is reduced to a point x_0 . We will say that s is based at x_0 .

If A is integrable, then one can use the reasoning of proposition A.1 to see that any A -sphere integrates to a unique sphere lying in the $\mathbf{s}$ -fiber over its base point, and based at an identity. This motivates the following definition of homotopy for A -spheres.

Definition C.2. A **homotopy** of A -spheres is a smooth family of A -spheres $s_u : TI^2 \rightarrow A, u \in I$ based at the same point $x_0 \in M$.

Definition C.3. Two spheres $s^0, s^1 : TI^2 \rightarrow A$ are said to be homotopic if there exists a homotopy of spheres $s^u : TI^2 \rightarrow A$ such that $s_i = s^i$, for $i = 0, 1$ $h = adt + bde + cdu \rightarrow A$ such that $a^{u=i}dt + b^{u=i}de = s^i$, for $i = 0, 1$.

We get this way an equivalence relation on the space of A -spheres.

Definition C.4. We call second homotopy group, denoted $\pi_2(A)$ the space of A -spheres up to homotopy.

The above definition for homotopy might not look so natural. Let us work a little bit and see why this notion is correct.

Proposition C.5. *Given a family (parametrized by $u \in I$) of spheres $s^u = a^u dt + b^u d\epsilon : TI^2 \rightarrow A$ based at a same point, there exists a unique $c : I^3 \rightarrow A$ such that $h : adt + bd\epsilon + cdu : TI^3 \rightarrow A$ is an algebroid morphism and $c|_{t,\epsilon=0} = 0$. Moreover, $c|_{t,\epsilon=1} = 0$*

Proof. Let us first show uniqueness: assume $h : adt + bd\epsilon + cdu : TI^3 \rightarrow A$ is an algebroid morphism, then one can extend b into a time-dependant section β of A chosen such that $\beta|_{t=0,1} = 0$, and let α be the solution of the evolution equation

$$\frac{d\alpha}{d\epsilon} - \frac{d\beta}{dt} = [\alpha, \beta]_A$$

with initial condition $\alpha|_{\epsilon=0} = 0$. We already know that necessarily $\alpha(\gamma^u(t, \epsilon)) = a(\epsilon, t, u)$ (see [6]). For the same reason, if we let θ be the solution of the evolution equation

$$\frac{d\theta}{d\epsilon} - \frac{d\beta}{du} = [\theta, \beta]_A,$$

with initial condition $\theta|_{\epsilon=0} = 0$, then necessarily θ satisfies $\theta_{\gamma^u(t, \epsilon)} = c(t, \epsilon, u)$.

To get existence, we have to make sure that if we let $c := \theta_{\gamma^u(t, \epsilon)}$, then $h : adt + bd\epsilon + cdu$ is indeed a Lie algebroid morphism. By construction, one already has the following relations:

$$\begin{aligned} \frac{d\alpha}{d\epsilon} - \frac{d\beta}{dt} &= [\alpha, \beta]_A \\ \frac{d\theta}{d\epsilon} - \frac{d\beta}{du} &= [\theta, \beta]_A \end{aligned}$$

and, as can be checked in local coordinates, it is enough to show that:

$$\frac{d\theta}{dt} - \frac{d\alpha}{du} = [\theta, \alpha]_A$$

as well. We will need this result for other purposes, so we stated it in a separate lemma (see below). There only remains to show that $\theta_{\gamma^{\epsilon(t)}|_{\epsilon=1}} = 0$. For this, one can make the same reasoning as above switching the roles of a and b , then invoke uniqueness. $\square$

Lemma C.6. *Let β be a smooth family of sections of an algebroid A with parameters $(\epsilon, t, u) \in I^3$ such that $\beta|_{t=0,1}$ vanishes. Denote α and θ the unique solutions of the evolution equations:*

$$\begin{aligned} \frac{d\alpha}{d\epsilon} - \frac{d\beta}{dt} &= [\alpha, \beta]_A, \\ \frac{d\theta}{d\epsilon} - \frac{d\beta}{du} &= [\theta, \beta]_A, \end{aligned}$$

with initial conditions $\alpha|_{\epsilon=0} = 0$ and $\theta|_{\epsilon=0} = 0$. Then the following assertions hold:

(i) θ is solution of:

$$\frac{d\theta}{dt} - \frac{d\alpha}{du} = [\theta, \alpha]_A,$$

(ii) θ vanishes whenever $t = 0, 1$.

Proof. Define a family of sections ϕ as the difference:

$$\phi := \frac{d\theta}{dt} - \frac{d\alpha}{du} - [\theta, \alpha]_A.$$

Clearly, $\phi|_{\epsilon=0} = 0$ since $\alpha|_{\epsilon=0}$ and $\theta|_{\epsilon=0}$ both vanish. On the other hand, a straightforward computation shows that ϕ also satisfies:

$$\frac{d\phi}{d\epsilon} = [\beta, \phi]_A.$$

The unique solution of such an equation with initial condition $\phi|_{\epsilon=0} = 0$ being $\phi = 0$, we have proved (i).

The fact that $\theta|_{t=0,1} = 0$ is in fact straightforward: if one sees the equation

$$\frac{d\theta}{d\epsilon} - \frac{d\beta}{du} = [\theta, \beta]_A,$$

as a family of equations with parameter t , clearly for $t = 0, 1$, since β vanishes, θ has to vanish as well, which leaves (ii) proved. $\square$

This way, we get an alternative definition for homotopies of A -spheres that should be more natural (though redundant).

Definition C.7. Two A -spheres $s^0, s^1 : TI^2 \rightarrow A$ are homotopic if there exists a Lie algebroid morphism $s : adt + bde + cdu : TI^3$ with $(adt + bde)|_{u=i} = s^i$ for $i = 0, 1$, and satisfying

- i) $a|_{\epsilon=0,1} = 0$ and $b|_{t=0,1} = 0$ (i.e., $adt + bde$ is an A -sphere for any $u \in I$).
- ii) $c|_{t,\epsilon=0,1} = 0$ (i.e., c vanishes whenever $t \in \{0, 1\}$ or $\epsilon \in \{0, 1\}$).

REFERENCES

- [1] O. Brahic and R. L. Fernandes, Poisson Fibrations and Fibered Symplectic Groupoids, Proceedings of the Conference on Poisson Geometry in Mathematics and Physics, Tokyo, 2006, AMS Contemporary Mathematics Series, 2008, no. 450, pages 41-60.
- [2] O. Brahic and R. L. Fernandes, *Integration of coupling Dirac structures*, in preparation.
- [3] A. Cannas da Silva, A. Weinstein, *Geometric models for noncommutative algebras*, Berkeley Math. Lecture Notes, 10, Amer. Math. Soc. (1999).
- [4] A.S. Cattaneo and G. Felder, Poisson sigma models and symplectic groupoids, in *Quantization of Singular Symplectic Quotients*, (ed. N. P. Landsman, M. Pflaum, M. Schlichenmeier), Progress in Mathematics **198** (2001), 41-73.
- [5] M. Crainic, Prequantization and Lie brackets, *J. Symplectic Geometry*, Vol.2, No.4, 579-602, 2005.
- [6] M. Crainic and R. L. Fernandes, Integrability of Lie brackets, *Ann. of Math. (2)* **157** (2003), 575-620.
- [7] M. Crainic and R. L. Fernandes, *Exotic Characteristic Classes of Lie Algebroids*, Quantum Field Theory and Noncommutative Geometry, Lecture Notes in Physics, Vol. 662. Eds. Carow-Watamura, Ursula; Maeda, Yoshiaki; Watamura, Satoshi, Springer-Verlag, Berlin, 2005.
- [8] M. Crainic and C. Zhu, *Integrability of Jacobi and Poisson structures*, Annales de l'institut Fourier, 57 no. 4 (2007), p. 1181-1216.
- [9] C. Ehresmann, Les connexions infinitésimales dans un espace fibré différentiable, *Séminaire Bourbaki*, Vol. 1, Exp. No. 24, 153-168, Soc. Math. France, Paris, 1995.
- [10] V. Ginzburg, Equivariant Poisson cohomology and a spectral sequence associated with a moment map, *Int. J. Math.*, **10** (1999), 977-1010.

- [11] V. Ginzburg, J. H. Lu, Poisson cohomology of Morita equivalent Poisson manifolds, *IMRN*, 10 (1992), 199-205.
- [12] P.J. Higgins, K. Mackenzie, *Algebraic constructions in the category of Lie algebroids*, J. Algebra, 129 (1990), 194-230.
- [13] K. Mackenzie, *Lie groupoids and Lie algebroids in differential geometry*, Cambridge Univ. Press (1987).
- [14] J. Mrcun and I. Moerdijk, *On integrability of infinitesimal actions*, Amer. J. Math. 124 (2002), 567-593.
- [15] Mackenzie and P. Xu P. *Integration of Lie bialgebroids*, Topology, Volume 39, Number 3, May 2000 , pp. 445-467(23)
- [16] J. Pradines, *Théorie de Lie pour les groupoïdes différentiables. Calcul différentiel dans la catégorie des groupoïdes infinitésimaux* C.R. Acad. Sci. Paris , **264** A (1967), 245-248.
- [17] H. Tseng and C. Zhu, *Integrating Lie algebroids via stacks*, Composition Mathematica 142 (2006), no.1, 251-270.
- [18] Y. Vorobjev, *Coupling tensors and Poisson geometry near a single symplectic leaf*, Banach Center Publ. **54** (2001), 249-274.
- [19] A. Wade, *Poisson fiber bundles and coupling Dirac structures*, Ann Glob Anal Geom (2008) 33, 207-217.
- [20] A. Weinstein, *Linearization of regular proper groupoids*, J. Inst. Math. Jussieu 1 (2002), no.3, 493-511.
- [21] P. Xu, *Poisson cohomology of regular Poisson manifolds*, Annales de l'institut Fourier, 42 no. 4 (1992), p. 967-988.

DEPART. DE MATEMÁTICA, INSTITUTO SUPERIOR TÉCNICO, 1049-001 LISBOA, PORTUGAL

E-mail address: brahic@math.ist.utl.pt